

#2311 Store at -20°C

Phospho-Synapsin (Ser9) Antibody

✓ 100 µl
(10 western blots)



Orders ■ 877-616-CELL (2355)
orders@cellsignal.com
Support ■ 877-678-TECH (8324)
info@cellsignal.com
Web ■ www.cellsignal.com

rev. 07/28/10

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Entrez-Gene ID #6853
Swiss-Prot Acc. #P17600

Applications	Species Cross-Reactivity*	Molecular Wt.	Source
W Endogenous	H, M, R	77 kDa	Rabbit**


Background: Synapsins, a group of at least five related members (synapsins Ia, Ib, IIa, IIb and IIIa), are abundant brain proteins essential for regulating neurotransmitter release (1,2). All synapsins contain a short amino-terminal domain that is highly conserved and phosphorylated by PKA or CaM kinase I (1). Phosphorylation of synapsin amino-terminal domain at serine 9 inhibits its binding to phospholipids and dissociates synapsins from synaptic vesicles (2).

Specificity/Sensitivity: Phospho-Synapsin (Ser9) Antibody detects endogenous levels of synapsin only when phosphorylated at serine 9. The antibody is expected to recognize phosphorylated serine 9 of all synapsin isoforms.

Source/Purification: Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding serine 9 of human synapsin I. Antibodies are purified by protein A and peptide affinity chromatography.

Background References:

- (1) Greengard, P. (1987) *Mol. Neurobiol.* 1, 81-119.
- (2) Hosaka, M. et al. (1999) *Neuron* 24, 377-387.



Western blot analysis of mouse brain homogenates, untreated or phosphatase-treated, using Phospho-Synapsin (Ser9) Antibody (upper) or Synapsin Antibody #2312 (lower).

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

*Species cross-reactivity is determined by western blot.

**Anti-rabbit secondary antibodies must be used to detect this antibody.

Recommended Antibody Dilutions:

Western blotting 1:1000

For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.