

#2526 Store at -20°C

Phospho-p53 (Ser33) Antibody

Small 100 μ l
 (10 western blots)



Orders ■ 877-616-CELL (2355)
 orders@cellsignal.com
Support ■ 877-678-TECH (8324)
 info@cellsignal.com
Web ■ www.cellsignal.com

rev. 04/01/10

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.


Entrez-Gene ID #7157
Swiss-Prot Acc. #P04637

Applications	Species Cross-Reactivity*	Molecular Wt.	Source
W, IHC-P Endogenous	H, (Mk)	53 kDa	Rabbit**

Background: The p53 tumor suppressor protein plays a major role in cellular response to DNA damage and other genomic aberrations. Activation of p53 can lead to either cell cycle arrest and DNA repair or apoptosis (1). p53 is phosphorylated at multiple sites *in vivo* and by several different protein kinases *in vitro* (2,3). DNA damage induces phosphorylation of p53 at Ser15 and Ser20 and leads to a reduced interaction between p53 and its negative regulator, the oncoprotein MDM2 (4). MDM2 inhibits p53 accumulation by targeting it for ubiquitination and proteasomal degradation (5,6). p53 can be phosphorylated by ATM, ATR and DNA-PK at Ser15 and Ser37. Phosphorylation impairs the ability of MDM2 to bind p53, promoting both the accumulation and activation of p53 in response to DNA damage (4,7). Chk2 and Chk1 can phosphorylate p53 at Ser20, enhancing its tetramerization, stability and activity (8,9). p53 is phosphorylated at Ser392 *in vivo* (10,11) and by CAK *in vitro* (11). Phosphorylation of p53 at Ser392 is increased in human tumors (12) and has been reported to influence the growth suppressor function, DNA binding and transcriptional activation of p53 (10,13,14). p53 is phosphorylated at Ser6 and Ser9 by CK1 δ and CK1 ϵ both *in vitro* and *in vivo* (13,15). Phosphorylation of p53 at Ser46 regulates the ability of p53 to induce apoptosis (16). Acetylation of p53 is mediated by p300 and CBP acetyltransferases. Inhibition of deacetylation suppressing MDM2 from recruiting HDAC1 complex by p19 (ARF) stabilizes p53. Acetylation appears to play a positive role in the accumulation of p53 protein in stress response (17). Following DNA damage, human p53 becomes acetylated at Lys382 (Lys379 in mouse) *in vivo* to enhance p53-DNA binding (18). Deacetylation of p53 occurs through interaction with the SIRT1 protein, a deacetylase that may be involved in cellular aging and the DNA damage response (19).

In vivo phosphorylation of p53 at Ser33 by CDK7/cyclin H and in response to UV-irradiation has been observed (17,18).

Specificity/Sensitivity: Phospho-p53 (Ser33) Antibody detects endogenous levels of p53 only when phosphorylated at serine 33.



Western blot analysis of extracts from HT29 cells treated with nocodazole (50 ng/ml, 24 h), calyculin A #9902 (100 nM, 10 min), or λ -Phosphatase NEB #P0753 (10,000 units/ml, 1 h), using Phospho-p53 (Ser33) Antibody (upper) or p53 (1C12) Mouse mAb #2524 (lower).

Source/Purification: Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser33 of human p53. Antibodies are purified by protein A and peptide affinity chromatography.

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

***Species cross-reactivity is determined by western blot.**

****Anti-rabbit secondary antibodies must be used to detect this antibody.**

Recommended Antibody Dilutions:

Western blotting	1:1000
Immunohistochemistry (Paraffin)	1:500
Unmasking buffer:	Citrate
Antibody diluent:	TBST-5%NGS

For application specific protocols please see the web page for this product at www.cellsignal.com.



Please visit www.cellsignal.com for a complete listing of recommended companion products.

Background References:


- Levin, A.J. (1997) *Cell* 88, 323-331.
- Meek, D.W. (1994) *Semin. Cancer Biol.* 5, 203-210.
- Milczarek, G.J. et al. (1997) *Life Sci.* 60, 1-11.
- Shieh, S.Y. et al. (1997) *Cell* 91, 325-334.
- Chehab, N.H. et al. (1999) *Proc. Natl. Acad. Sci. USA* 96, 13777-13782.
- Honda, R. et al. (1997) *FEBS Lett.* 420, 25-27.
- Tibbetts, R.S. et al. (1999) *Genes Dev.* 13, 152-157.
- Shieh, S.Y. et al. (1999) *EMBO J.* 18, 1815-1823.
- Hirao, A. et al. (2000) *Science* 287, 1824-1827.
- Hao, M. et al. (1996) *J. Biol. Chem.* 271, 29380-29385.
- Lu, H. et al. (1997) *Mol. Cell. Biol.* 17, 5923-5934.
- Ullrich, S.J. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90, 5954-5958.
- Kohn, K.W. (1999) *Mol. Biol. Cell* 10, 2703-2734.
- Lohrum, M. and Scheidtmann, K.H. (1996) *Oncogene* 13, 2527-2539.
- Knippsschild, U. et al. (1997) *Oncogene* 15, 1727-1736.
- Oda, K. et al. (2000) *Cell* 102, 849-862.
- Ito, A. et al. (2001) *EMBO J.* 20, 1331-1340.
- Sakaguchi, K. et al. (1998) *Genes Dev.* 12, 2831-2841.
- Solomon, J.M. et al. (2006) *Mol. Cell. Biol.* 26, 28-38.
- Ko, L. J. et al. (1997) *Mol. Cell. Biol.* 17, 7220-7229.
- Latonen, L. et al. (2001) *Oncogene* 20, 6784-6793.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.


Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
 Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.



Immunohistochemical analysis of paraffin-embedded human lung dysplasia of alveolar cells, using Phospho-p53 (Ser33) Antibody in the presence of control peptide (left) or antigen specific peptide (right),



Immunohistochemical analysis of paraffin-embedded human breast carcinoma, using Phospho-p53 (Ser33) Antibody.



Immunohistochemical analysis of paraffin-embedded human colon carcinoma, showing nuclear localization, using Phospho-p53 (Ser33) Antibody.