

#2543
Store at -20C

Phospho-cdc2 (Thr14) Antibody

**Cell Signaling**
TECHNOLOGY®**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

| Applications: | Reactivity: | MW (kDa): | Source: | UniProt ID: | Entrez-Gene Id: |
|---------------|-------------|-----------|---------|-------------|-----------------|
| WB | H Hm Mk | 34 | Rabbit | #P06493 | 983 |

| | | |
|--|--|-----------------|
| Product Usage Information | Application | Dilution |
| | Western Blotting | 1:1000 |
| Storage | Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody. | |
| Specificity / Sensitivity | Phospho-cdc2 (Thr14) Antibody detects endogenous levels of cdc2 (CDK1) only when phosphorylated at Thr14. Based on sequence similarity, the antibody may cross-react with CDK2 and CDK3. | |
| Species predicted to react based on 100% sequence homology: | Mouse, Rat, D. melanogaster, Xenopus, Bovine | |
| Source / Purification | Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Thr14 of human cdc2. Antibodies are purified by protein A and peptide affinity chromatography. | |
| Background | The entry of eukaryotic cells into mitosis is regulated by cdc2 kinase activation, a process controlled at several steps including cyclin binding and phosphorylation of cdc2 at Thr161 (1). However, the critical regulatory step in activating cdc2 during progression into mitosis appears to be dephosphorylation of cdc2 at Thr14 and Tyr15 (2). Phosphorylation at Thr14 and Tyr15, resulting in inhibition of cdc2, can be carried out by Wee1 and Myt1 protein kinases (3,4). The cdc25 phosphatase may be responsible for removal of phosphates at Thr14 and Tyr15 and subsequent activation of cdc2 (1,5). | |
| Background References | <ol style="list-style-type: none"> 1. Atherton-Fessler, S. et al. (1994) <i>Mol Biol Cell</i> 5, 989-1001. 2. Norbury, C. et al. (1991) <i>EMBO J</i> 10, 3321-9. 3. McGowan, C.H. and Russell, P. (1993) <i>EMBO J</i> 12, 75-85. 4. Wells, N.J. et al. (1999) <i>J Cell Sci</i> 112 (Pt 19), 3361-71. 5. Hunter, T. (1995) <i>Cell</i> 80, 225-36. | |

| | |
|-------------------------------|---|
| Species Reactivity | Species reactivity is determined by testing in at least one approved application (e.g., western blot). |
| Western Blot Buffer | IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight. |
| Applications Key | WB: Western Blotting |
| Cross-Reactivity Key | H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected |
| Trademarks and Patents | Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information. |
| Limited Uses | Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect. |

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.