

#2674 Store at -20°C

Phospho-53BP1 (Ser25/29) Antibody

✓ 100 µl
(10 western blots)



Orders ■ 877-616-CELL (2355)
orders@cellsignal.com
Support ■ 877-678-TECH (8324)
info@cellsignal.com
Web ■ www.cellsignal.com

rev. 03/30/10

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.


Entrez-Gene ID #7158
Swiss-Prot Acc. #Q12888

Applications	Species Cross-Reactivity*	Molecular Wt.	Source
W Endogenous	H, Mk	450 kDa	Rabbit**

Background: p53-binding protein 1 (53BP1) was originally identified as a p53 binding partner that could enhance the transcriptional activity of p53 (1,2). 53BP1 consists of two BRCA1 carboxy-terminal (BRCT) domains that allow for binding to p53 and a separate domain responsible for binding to phosphorylated histone H2A.X (3). 53BP1 rapidly translocates to nuclear foci following treatment of cells with ionizing radiation (IR) or radiomimetic agents that cause DNA double strand breaks (DSBs) (4,5). Because of this localization to DSBs and homology to the yeast protein Rad9, a role for 53BP1 in DSB repair has been proposed. Recruitment of 53BP1 to sites of DNA damage has been demonstrated to be independent of ATM, NBS1, and DNA-PK (4) and retention of 53BP1 at DNA breaks requires phosphorylated H2A.X (6). In cells lacking 53BP1, phosphorylation of ATM substrates is reduced, suggesting that 53BP1 is upstream of ATM (7). In response to IR, phosphorylation of 53BP1 at serines 6, 25, 29, and 784 by ATM has been demonstrated, but phosphorylation at these sites is not required for localization of 53BP1 to sites of DSBs (6).

Specificity/Sensitivity: Phospho-53BP1 (Ser25/29) Antibody detects endogenous levels of 53BP1 only when phosphorylated at serine 25/29.

Source/Purification: Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser25/29 of human 53BP1. Antibodies are purified by protein A and peptide affinity chromatography.



Western blot analysis of extracts from 293 cells, untreated or UV-treated (50 µJ for 2 hours), using Phospho-53BP1 (Ser25/29) Antibody (upper) or 53BP1 Antibody #4937 (lower).

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

*Species cross-reactivity is determined by western blot.

**Anti-rabbit secondary antibodies must be used to detect this antibody.

Recommended Antibody Dilutions:

Western Blotting 1:1000

For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

Background References:

- (1) Iwabuchi, K. et al. (1994) *Proc Natl Acad Sci U S A* 91, 6098–6102.
- (2) Iwabuchi, K. et al. (1998) *J Biol Chem* 273, 26061–26068.
- (3) Mochan, T.A. et al. (2004) *DNA Repair (Amst)* 3, 945–952.
- (4) Schultz, L.B. et al. (2000) *J Cell Biol* 151, 1381–1390.
- (5) Anderson, L. et al. (2001) *Mol Cell Biol* 21, 1719–1729.
- (6) Ward, I.M. et al. (2003) *J Biol Chem* 278, 19579–19582.
- (7) DiTullio, R.A. et al. (2002) *Nat Cell Biol* 4, 998–1002.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

© 2010 Cell Signaling Technology, Inc.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
 Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.