

#2717 Store at -20°C

Phospho-Zap-70 (Tyr319)/Syk (Tyr352) (65E4) Rabbit mAb

100 µl
 (10 western blots)



Orders ■ 877-616-CELL (2355)
 orders@cellsignal.com
Support ■ 877-678-TECH (8324)
 info@cellsignal.com
Web ■ www.cellsignal.com

rev. 08/23/10

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Entrez-Gene ID # 7535
Swiss-Prot Acc. # P43403

Applications	Species Cross-Reactivity*	Molecular Wt.	Isotype
W	H, M	70, 72 kDa	Rabbit IgG**
Endogenous			


Background: Zap-70, a Syk family protein tyrosine kinase expressed in T and NK cells, plays a critical role in mediating T cell activation in response to T cell receptor (TCR) engagement (1). Following TCR engagement, Zap-70 is rapidly phosphorylated on several tyrosine residues, presumably by two mechanisms: autophosphorylation and transphosphorylation by the Src family tyrosine kinase, Lck (2–6). Tyrosine phosphorylation of Zap-70 correlates with its increased kinase activity and downstream signaling events. In patients with chronic lymphocytic leukemia (CLL), total Zap-70 expression was shown to be correlated with disease progression and survival (7,8).

Specificity/Sensitivity: Phospho-Zap-70 (Tyr319)/Syk (Tyr352) (65E4) Rabbit mAb detects endogenous levels of Zap-70 only when phosphorylated at Tyr319. It cross-reacts with endogenous levels of Syk when phosphorylated at Tyr352.

Source/Purification: Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr319 of human Zap-70.

Background References:

- (1) Chu, D.H. et al. (1998) *Immunol. Rev.* 165, 167–180.
- (2) Iwashima, M. et al. (1994) *Science* 263, 1136–1139.
- (3) Neumeister, E.N. et al. (1995) *Mol. Cell. Biol.* 15, 3171–3178.
- (4) Chan, A.C. et al. (1995) *EMBO J.* 14, 2499–2508.
- (5) Williams, B.L. et al. (1999) *EMBO J.* 18, 1832–1844.
- (6) Di Bartolo, V. et al. (1999) *J. Biol. Chem.* 274, 6285–6294.
- (7) Wiestner, A. et al. (2003) *Blood* 101, 4944–4951.
- (8) Crespo, M. et al. (2003) *N. Engl. J. Med.* 348, 1764–1775.



Western blot analysis of extracts from Jurkat cells treated with hydrogen peroxide (2mM for 2 minutes) or with λ phosphatase and extracts from Ramos cells treated with anti-human IgM (12 micrograms/ml for 2 minutes) using Phospho-Zap-70 (Tyr319)/Syk (Tyr352) (65E4) Rabbit mAb.

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

*Species cross-reactivity is determined by western blot.

**Anti-rabbit secondary antibodies must be used to detect this antibody.

Recommended Antibody Dilutions:

Western Blotting 1:1000

For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

© 2010 Cell Signaling Technology, Inc. Rabbit monoclonal antibody is produced under license (granting certain rights including those under U. S. Patents No. 5,675,063 and 7,429,487) from Epitomics, Inc.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
 Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.