

Normal Rabbit IgG

✓ 250 µl

Orders ■ 877-616-CELL (2355)
orders@cellsignaling.com

Support ■ 877-678-TECH (8324)
info@cellsignaling.com

Web ■ www.cellsignaling.com


New 03/08

This product is for *in vitro* research use only and is not intended for use in humans or animals.

Concentration	Applications	Source	Isotype
1 mg/ml	ChIP	Rabbit	IgG

Description: Normal Rabbit IgG is an unconjugated rabbit polyclonal antibody that is routinely used as a non-specific IgG control in chromatin immunoprecipitation using our SimpleChIP™ Enzymatic Chromatin IP Kits #9002 and #9003.


Background: Normal Rabbit IgG is an isotype control antibody, which is used to estimate the non-specific binding of target primary antibodies due to Fc receptor binding or other protein-protein interactions. An isotype control antibody should have the same immunoglobulin type and be used at the same concentration as the test antibody.



Chromatin immunoprecipitations were performed using digested chromatin from HeLa cells and the indicated antibodies. Purified DNA was analyzed by quantitative Real-Time PCR, using primers specific for the RPL30 gene, the MYOD gene or the heterochromatic α satellite repeat element. The relative abundance of each DNA sequence enriched by Normal Rabbit IgG #2729 (red) is compared to the amount of the same DNA sequence enriched by the protein-specific immunoprecipitations.

Source/Purification: The purified antibody is not directed against any known antigen. It was isolated from naive rabbit sera and prepared by Protein A chromatography.

Directions for Use: Important! Dilute this control antibody to the same concentration (not dilution) as the specific test antibody in the chromatin immunoprecipitation. This is typically 1 µl (1 µg) to 5 µl (5 µg) of control antibody for one immunoprecipitation. Higher background signal may result if an excessive amount of rabbit IgG isotype control is used.



Chromatin immunoprecipitations were performed using digested chromatin from HeLa cells and either Histone H3 Antibody #2650 (lane 2), Rpb1 CTD (4H8) Mouse mAb #2629 (lane 3), Di-Methyl Histone H3 (Lys9) Antibody #9753 (lane 4) or Normal Rabbit IgG #2729 (lane 5). Purified DNA was analyzed by standard PCR methods using primers specific for the RPL30 gene, the MYOD gene or the heterochromatic α satellite repeat element. PCR products were observed for each primer set in the input sample (lane 1) and various protein-specific immunoprecipitations but no PCR products were observed with immunoprecipitation using Normal Rabbit IgG #2729 (lane 5).

Storage: Supplied in 10 mM HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. *Do not aliquot the antibody.*

Recommended Antibody Dilutions:

Chromatin IP:
Match IgG concentration of test antibody

Companion Products:

SimpleChIP™ Enzymatic Chromatin IP Kit (Agarose Beads) #9002
SimpleChIP™ Enzymatic Chromatin IP Kit (Magnetic Beads) #9003
ChIP-Grade Protein G Magnetic Beads #9006
ChIP-Grade Protein G Agarose Beads #9007
Di-Methyl-Histone H3 (Lys4) (C64G9) Rabbit mAb #9725
Di-Methyl-Histone H3 (Lys4) Antibody #9726
Tri-Methyl-Histone H3 (Lys4) (C42D8) Rabbit mAb #9751
Di-Methyl-Histone H3 (Lys9) Antibody #9753
Pan-Methyl-Histone H3 (Lys9) Antibody #4069
Tri-Methyl-Histone H3 (Lys27) Antibody #9756
Acetyl-Histone H3 (Lys9) (C5B11) Rabbit mAb #9649
Acetyl-Histone H3 (Lys9/Lys14) Antibody #9677
Rpb1 CTD (4H8) Mouse mAb #2629