

#2761 Store at -20°C

Numb (C44B4) Rabbit mAb

✓ 100 µl
(10 western blots)

Orders ■ 877-616-CELL (2355)
orders@cellsignaling.com
Support ■ 877-678-TECH (8324)
info@cellsignaling.com
Web ■ www.cellsignaling.com

rev. 08/12/10


This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Applications	Species Cross-Reactivity*	Molecular Wt.	Isotype
W, IP, IF-IC, F Endogenous	H	72, 74 kDa	Rabbit IgG**


Background: Numb contains an amino-terminal phosphotyrosine-binding (PTB) domain and carboxy-terminal endocytic binding motifs for α -adaptin and EH (Eps15 homology) domain-containing proteins, indicating a role in endocytosis (1,2). There are four mammalian Numb splicing isoforms that are differentially expressed and may have distinct functions (3-5). Numb acts as a negative regulator of Notch signaling by promoting ubiquitination and degradation of Notch (6). The protein is asymmetrically segregated into one daughter cell during cell division; therefore the two daughter cells have different responses to Notch signaling and have different cell fates (7,8). The localization of Numb can also be regulated by G-protein coupled receptor (GPCR) and PKC signaling (9).

Specificity/Sensitivity: Numb (C44B4) Rabbit mAb detects endogenous levels of Numb protein.


Source/Purification: Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Phe482 of human Numb protein.



Western blot analysis of extracts from HeLa and SK-OV-3 cells using Numb (C44B4) Rabbit mAb.



Flow cytometric analysis of A204 cells using Numb (C44B4) Rabbit mAb (blue) compared to a nonspecific negative control antibody (red).



Confocal immunofluorescent analysis of HeLa cells using Numb (C44B4) Rabbit mAb (green). Actin filaments have been labeled with DY-554 phalloidin (red). Blue pseudocolor = DRAQ5 (fluorescent DNA dye).

Entrez-Gene ID #8650
Swiss-Prot Acc. #P49757

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

*Species cross-reactivity is determined by western blot.

**Anti-rabbit secondary antibodies must be used to detect this antibody.

Recommended Antibody Dilutions:

Western Blotting	1:1000
Immunoprecipitation	1:50
Immunofluorescence (IF-IC)	1:100
Flow Cytometry	1:200

For application specific protocols please see the web page for this product at www.cellsignaling.com.

Please visit www.cellsignaling.com for a complete listing of recommended companion products.

Background References:

- (1) Berndt, D. et al. (2002) *Dev. Cell* 3, 221-231.
- (2) Santolini, E. et al. (2000) *J. Cell Biol.* 151, 1345-1352.
- (3) Dho, S.E. et al. (1999) *J. Biol. Chem.* 274, 33097-33104.
- (4) Verdi, J.M. et al. (1999) *Proc. Natl. Acad. Sci. USA* 96, 10472-10476.
- (5) Toriya, M. et al. (2006) *Dev. Neurosci.* 28, 142-155.
- (6) McGill, M.A. and McGlade, C.J. (2003) *J. Biol. Chem.* 278, 23196-23203.
- (7) Verdi, J.M. et al. (1996) *Curr. Biol.* 6, 1134-1145.
- (8) Reugels, A.M. et al. (2006) *Dev. Dyn.* 235, 934-948.
- (9) Dho, S.E. et al. (2006) *Mol. Biol. Cell.*, Epub Ahead of Print.

DRAQ5® is a registered trademark of Biostatus Limited.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
 Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.