

#2924 Store at -20°C

# Cyclin D2 Antibody

✓ 100 µl  
(10 western blots)



**Orders** ■ 877-616-CELL (2355)  
orders@cellsignaling.com  
**Support** ■ 877-678-TECH (8324)  
info@cellsignaling.com  
**Web** ■ www.cellsignaling.com

rev. 04/05/10

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

**Entrez-Gene ID** #894  
**Swiss-Prot Acc.** #P30279

Applications W Endogenous	Species Cross-Reactivity*		Molecular Wt. 30 kDa	Source Rabbit**
	H	M		


**Background:** Activity of the cyclin-dependent kinases CDK4 and CDK6 is regulated by T-loop phosphorylation, by the abundance of their cyclin partners (the D-type cyclins), and by association with CDK inhibitors of the Cip/Kip or INK family of proteins (1). The inactive ternary complex of cyclin D/CDK4 and p27 Kip1 requires extracellular mitogenic stimuli for the release and degradation of p27 concomitant with a rise in cyclin D levels to effect progression through the restriction point and pRb-dependent entry into S-phase (2). The active complex of cyclin D/CDK4 targets the retinoblastoma protein for phosphorylation, allowing the release of E2F transcription factors that activate G1/S-phase gene expression (3). Levels of cyclin D protein drop upon withdrawal of growth factors through downregulation of its protein expression and through phosphorylation-dependent degradation (4).

**Specificity/Sensitivity:** Cyclin D2 Antibody detects endogenous levels of cyclin D2. It does not cross-react with other family members at physiological levels.

**Source/Purification:** Polyclonal antibodies are produced by immunizing animals with a synthetic peptide to corresponding to residues near the amino terminus of cyclin D2. Antibodies are purified by protein A and peptide affinity chromatography.

**Background References:**

- (1) Hirai, H. et al. (1995) *Mol. Cell. Biol.* 15, 2672–2681.
- (2) Sherr, C.J. (1996) *Science* 274, 1672–1677.
- (3) Lukas, J. et al. (1996) *Mol. Cell. Biol.* 16, 6917–6925.
- (4) Diehl, J.A. et al. (1997) *Genes Dev.* 11, 957–972.



Western blot analysis of extracts from Raji cells, using Cyclin D2 Antibody.

**Storage:** Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

\*Species cross-reactivity is determined by western blot.

\*\*Anti-rabbit secondary antibodies must be used to detect this antibody.

**Recommended Antibody Dilutions:**  
Western blotting 1:1000

For application specific protocols please see the web page for this product at [www.cellsignaling.com](http://www.cellsignaling.com).

Please visit [www.cellsignaling.com](http://www.cellsignaling.com) for a complete listing of recommended companion products.

**IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.**

© 2010 Cell Signaling Technology, Inc.

**Applications Key:** W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide  
**Species Cross-Reactivity Key:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine  
 Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.