

#2965 Store at -20°C

# Phospho-Akt (Thr308) (C31E5E) Rabbit mAb

- Small 100 µl (10 western blots)
- Large 300 µl (30 western blots)

rev. 02/23/10



**Orders** ■ 877-616-CELL (2355)  
orders@cellsignal.com

**Support** ■ 877-678-TECH (8324)  
info@cellsignal.com

**Web** ■ www.cellsignal.com


This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Applications	Species Cross-Reactivity*	Molecular Wt.	Isotype
W, IF-IC, F Endogenous	H, M, R, Hm, Mk	60 kDa	Rabbit IgG**

**Background:** Akt, also referred to as PKB or Rac, plays a critical role in controlling survival and apoptosis (1-3). This protein kinase is activated by insulin and various growth and survival factors to function in a wortmannin-sensitive pathway involving PI3 kinase (2,3). Akt is activated by phospholipid binding and activation loop phosphorylation at Thr308 by PDK1 (4) and by phosphorylation within the carboxy terminus at Ser473. The previously elusive PDK2 responsible for phosphorylation of Akt at Ser473 has been identified as mammalian target of rapamycin (mTOR) in a rapamycin-insensitive complex with rictor and Sin1 (5,6). Akt promotes cell survival by inhibiting apoptosis by phosphorylating and inactivating several targets, including Bad (7), forkhead transcription factors (8), c-Raf (9) and caspase-9. PTEN phosphatase is a major negative regulator of the PI3 kinase/Akt signaling pathway (10). LY294002 is a specific PI3 kinase inhibitor (11).

Another essential Akt function is the regulation of glycogen synthesis through phosphorylation and inactivation of GSK-3α and β (12,13). Akt may also play a role in insulin stimulation of glucose transport (12).

In addition to its role in survival and glycogen synthesis, Akt is involved in cell cycle regulation by preventing GSK-3β mediated phosphorylation and degradation of cyclin D1 (14) and by negatively regulating the cyclin dependent kinase inhibitors p27 Kip (15) and p21 Waf1 (16). Akt also plays a critical role in cell growth by directly phosphorylating mTOR in a rapamycin-sensitive complex containing raptor (17). More importantly, Akt phosphorylates and inactivates tuberlin (TSC2), an inhibitor of mTOR within the mTOR-raptor complex (18). Inhibition of mTOR stops the protein synthesis machinery due to inactivation of its effector, p70 S6 kinase and activation of the eukaryotic initiation factor 4E binding protein 1 (4E-EP1), an inhibitor of translation (18,19).



Western blot analysis of extracts from NIH/3T3 and Jurkat cells, untreated, PDGF-treated or LY294002-treated as indicated, using Phospho-Akt (Thr308) (C31E5E) Rabbit mAb (upper) or Akt (pan) (C67E7) Rabbit mAb #4691 (lower).

**Specificity/Sensitivity:** Phospho-Akt (Thr308) (C31E5E) Rabbit mAb detects endogenous levels of Akt only when phosphorylated at Thr308.

**Source/Purification:** Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues around Thr308 of mouse Akt.

Entrez-Gene ID #207  
Swiss-Prot Acc. #P31749

**Storage:** Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

**\*Species cross-reactivity is determined by western blot.**  
**\*\*Anti-rabbit secondary antibodies must be used to detect this antibody.**


**Recommended Antibody Dilutions:**

Western blotting	1:1000
Immunofluorescence (IF-IC)	1:800
Flow Cytometry	1:100

For application specific protocols please see the web page for this product at [www.cellsignal.com](http://www.cellsignal.com).

Please visit [www.cellsignal.com](http://www.cellsignal.com) for a complete listing of recommended companion products.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc.  
DRAQ5® is a registered trademark of Biostatus Limited.





**IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.**

**Applications Key:** W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide


**Species Cross-Reactivity Key:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine

Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.

© 2010 Cell Signaling Technology, Inc. Rabbit monoclonal antibody is produced under license (granting certain rights including those under U. S. Patent No. 5,675,063) from EpiTomics, Inc.



Confocal immunofluorescent analysis of C2C12 cells, insulin-treated (left) or LY294002-treated (right), using Phospho-Akt (Thr308) (C31E5E) Rabbit mAb (green). Actin filaments have been labeled with Alexa Fluor<sup>®</sup> 555 phalloidin (red). Blue pseudocolor = DRAQ5<sup>®</sup> #4084 (fluorescent DNA dye).



Flow cytometric analysis of Jurkat cells, untreated (green) or LY294002, Wortmannin and U0126-treated (blue) using Phospho-Akt (Thr308) (C31E5E) Rabbit mAb.

#### Background References:

- (1) Franke, T.F. et al. (1997) *Cell* 88, 435–7.
- (2) Burgering, B.M. and Coffey, P.J. (1995) *Nature* 376, 599–602.
- (3) Franke, T.F. et al. (1995) *Cell* 81, 727–36.
- (4) Alessi, D.R. et al. (1996) *EMBO J* 15, 6541–51.
- (5) Sarbassov, D.D. et al. (2005) *Science* 307, 1098–101.
- (6) Jacinto, E. et al. (2006) *Cell* 127, 125–37.
- (7) Cardone, M.H. et al. (1998) *Science* 282, 1318–21.
- (8) Brunet, A. et al. (1999) *Cell* 96, 857–68.
- (9) Zimmermann, S. and Moelling, K. (1999) *Science* 286, 1741–4.
- (10) Cantley, L.C. and Neel, B.G. (1999) *Proc Natl Acad Sci USA* 96, 4240–5.
- (11) Vlahos, C.J. et al. (1994) *J Biol Chem* 269, 5241–8.
- (12) Hajdich, E. et al. (2001) *FEBS Lett* 492, 199–203.
- (13) Cross, D.A. et al. (1995) *Nature* 378, 785–9.
- (14) Diehl, J.A. et al. (1998) *Genes Dev* 12, 3499–511.
- (15) Gesbert, F. et al. (2000) *J Biol Chem* 275, 39223–30.
- (16) Zhou, B.P. et al. (2001) *Nat Cell Biol* 3, 245–52.
- (17) Navé, B.T. et al. (1999) *Biochem J* 344 Pt 2, 427–31.
- (18) Inoki, K. et al. (2002) *Nat Cell Biol* 4, 648–57.
- (19) Manning, B.D. et al. (2002) *Mol Cell* 10, 151–62.