

#3387 Store at -20°C

Cdc6 (C42F7) Rabbit mAb



✓ 100 µl
(10 Western mini-blot)

Orders ■ 877-616-CELL (2355)
orders@cellsignaling.com
Support ■ 877-678-TECH (8324)
info@cellsignaling.com
Web ■ www.cellsignaling.com

New 11/08

This product is for *in vitro* research use only and is not intended for use in humans or animals.
This product is not intended for use as a therapeutic or in diagnostic procedures.

Entrez-Gene ID #990
Swiss-Prot Acc. #Q99741


Applications	Species Cross-Reactivity*	Molecular Wt.	Isotype
W Endogenous	H, M, R, Mk, Hm	62 kDa	Rabbit IgG**

Background: The initiation of DNA replication in mammalian cells is a highly coordinated process that ensures duplication of the genome only once per cell division cycle. Origins of replication are dispersed throughout the genome, and their activities are regulated via the sequential binding of pre-replication and replication factors. The origin recognition complex (ORC) is thought to be bound to chromatin throughout the cell cycle (1,2). The pre-replication complex (Pre-RC) forms in late mitosis/early G1 phase beginning with the binding of CDT1 and cdc6 to the origin, which allows binding of the heterohexameric MCM2-7 complex. The MCM complex is thought to be the replicative helicase, and formation of the pre-RC is referred to as chromatin licensing. Subsequent initiation of DNA replication requires the activation of the S-phase promoting kinases cdk2 and cdc7. Cdc7, which is active only in complex with its regulatory subunit dbf4, phosphorylates MCM proteins bound to chromatin and allows binding of the replication factor cdc45 and DNA polymerase (3,4).

Replication licensing is controlled in part by the degradation of cdc6 in quiescent cells. Phosphorylation of cdc6 by cdk2 prevents its degradation, allowing pre-replication complexes to form (5). Cdc6 has recently been shown to play an important role in the intra-S-phase p21 Waf1/Cip1-dependent DNA damage response (6,7). Both cdc6 and CDT1 are degraded by the ubiquitin proteasome pathway in response to DNA damage associated with re-replication (8).

Specificity/Sensitivity: Cdc6 (C42F7) Rabbit mAb detects endogenous levels of total cdc6 protein.

Source/Purification: Monoclonal antibody is produced by immunizing animals with a synthetic peptide (KLH-coupled) corresponding to residues near the amino terminus of human cdc6.



Western blot analysis of extracts from various cell types using Cdc6 (C42F7) Rabbit mAb.

Background References:

- (1) Okuno, Y. et al. (2001) *EMBO J* 20, 4263–77.
- (2) McNairn, A.J. et al. (2005) *Exp Cell Res* 308, 345–56.
- (3) Bell, S.P. and Dutta, A. (2002) *Annu Rev Biochem* 71, 333–74.
- (4) Tsuji, T. et al. (2006) *Mol Biol Cell* 17, 4459–72.
- (5) Mailand, N. and Diffley, J.F. (2005) *Cell* 122, 915–26.
- (6) Kan, Q. et al. (2008) *J Biol Chem* 283, 17864–72.
- (7) Kan, Q. et al. (2008) *Proc Natl Acad Sci USA* 105, 4757–62.
- (8) Hall, J.R. et al. (2008) *J Biol Chem* 283, 25356–63.

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

*Species cross-reactivity is determined by Western blot.

**Anti-rabbit secondary antibodies must be used to detect this antibody.

Recommended Antibody Dilutions:

Western blotting 1:1000

For application specific protocols please see the web page for this product at www.cellsignaling.com.

Companion Products:

- MCM2 Antibody #4007
- MCM3 Antibody #4012
- MCM7 Antibody #4018
- ORC2 (3G6) Rat mAb #4736
- ORC6 (3A4) Rat mAb #4737
- Anti-rabbit IgG, HRP-linked Antibody #7074
- Phototope®-HRP Western Blot Detection System, Anti-rabbit IgG, HRP-linked Antibody #7071
- Prestained Protein Marker, Broad Range (Premixed Format) #7720
- Biotinylated Protein Ladder Detection Pack #7727
- 20X LumiGLO® Reagent and 20X Peroxide #7003

Please visit www.cellsignaling.com for a complete listing of recommended companion products.

IMPORTANT: For Western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebra fish B—bovine
 Dg—dog Pg—pig Sc—S. cerevisiae All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.