

#3670 Store at -20°C

GFAP (GA5) Mouse mAb

100 µl
 (10 Western mini-blot)



Orders ■ 877-616-CELL (2355)
 orders@cellsignal.com
Support ■ 877-678-TECH (8324)
 info@cellsignal.com
Web ■ www.cellsignal.com

rev. 01/14/09


This product is for *in vitro* research use only and is not intended for use in humans or animals.
 This product is not intended for use as a therapeutic or in diagnostic procedures.

Applications	Species Cross-Reactivity*	Molecular Wt.	Source	Isotype
W, IP, IHC-P, IF-F Endogenous	H, M, R	50 kDa	Mouse	IgG1


Background: The cytoskeleton consists of three types of cytosolic fibers: microfilaments (actin filaments), intermediate filaments and microtubules. Major types of intermediate filaments are distinguished and expressed in particular cell types: cytokeratins (epithelial cells), glial fibrillary acidic protein, GFAP (glial cells), desmin (skeletal, visceral and certain vascular smooth muscle cells), vimentin (mesenchyme origin) and neurofilaments (neurons). GFAP and vimentin form intermediate filaments in astroglial cells and modulate their motility and shape (1). In particular, vimentin filaments are present at early developmental stages, while GFAP filaments are characteristic of differentiated and mature brain astrocytes. Thus, GFAP is commonly used as a marker for intracranial and intraspinal tumors arising from astrocytes (2). In addition, GFAP intermediate filaments are also present in non-myelin forming Schwann cells in the peripheral nervous system (3).

Specificity/Sensitivity: GFAP (GA5) Mouse mAb detects endogenous levels of total GFAP protein.

Source/Purification: Monoclonal antibody is produced by immunizing mice with native GFAP purified from pig spinal cord.



Western blot analysis of extracts from mouse and rat brain, using GFAP (GA5) Mouse mAb.



Immunohistochemical staining analysis of paraffin-embedded human medulloblastoma, using GFAP (GA5) Mouse mAb.

Background References:

- (1) Eng, L.F. et al. (2000) *Neurochem. Res.* 25, 1439–51.
- (2) Goebel, H.H. et al. (1987) *Acta. Histochem. Suppl.* 34, 81–93.
- (3) Jessen, K.R. et al. (1990) *Development* 109, 91–103.

Entrez-Gene ID #2670
 Swiss-Prot Acc. #P14136

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

***Species cross-reactivity is determined by Western blot.**

****Anti-mouse secondary antibodies must be used to detect this antibody.**

Recommended Antibody Dilutions:


Western blotting	1:1000
Immunoprecipitation	1:50
Immunohistochemistry (Paraffin)	1:50
IHC protocol: Unmasking buffer/Antibody diluent Citrate/TBST-5%NGS	
Immunofluorescence (IF-F)	1:300

For application specific protocols please see the web page for this product at www.cellsignal.com.

Companion Products:

- α/β-Tubulin Antibody #2148
- Phototope®-HRP Western Blot Detection System, Anti-mouse IgG, HRP-linked Antibody #7072
- Anti-mouse IgG, HRP-linked Antibody #7076
- Prestained Protein Marker, Broad Range (Premixed Format) #7720
- Biotinylated Protein Ladder Detection Pack #7727
- 20X LumiGLO® Reagent and 20X Peroxide #7003

Please visit www.cellsignal.com for a complete listing of recommended companion products.



Confocal immunofluorescence image of rat hippocampus labeled with GFAP (GA5) Mouse mAb (red), Phospho-S6 Ribosomal Protein (Ser235/236) (2F9) Rabbit mAb (Alexa Fluor® 488 Conjugate) #4854 (green), and CREB (48H2) Rabbit mAb #9197 (blue).

IMPORTANT: For Western blots, incubate membrane with diluted antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebra fish B—bovine
 Dg—dog Pg—pig Sc—S. cerevisiae All—all species expected Species enclosed in parentheses are predicted to react based on 100% sequence homology.