

#3765 Store at **-20°C**

Phospho-ASK1 (Thr845) Antibody

100 µl
 (10 western blots)



Orders ■ 877-616-CELL (2355)
 orders@cellsignal.com
Support ■ 877-678-TECH (8324)
 info@cellsignal.com
Web ■ www.cellsignal.com

rev. 02/01/10

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Applications W Transfected	Species Cross-Reactivity* H, (M)	Molecular Wt. 155 kDa	Source Rabbit**
----------------------------------	-------------------------------------	--------------------------	--------------------


Background: Apoptosis signal-regulating kinase 1 (ASK1), a MAP kinase kinase kinase, plays essential roles in stress-induced apoptosis (1,2). ASK1 is activated in response to a variety of stress-related stimuli through distinct mechanisms and activates MKK4 and MKK3, which in turn activate JNK and p38 (3). Overexpression of ASK1 activates JNK and p38 and induces apoptosis in several cell types through signals involving the mitochondrial cell death pathway. Embryonic fibroblasts or primary neurons derived from ASK1^{-/-} mice are resistant to stress-induced JNK and p38 activation and cell death (4,5). Phosphorylation at Ser967 is essential for ASK1 association with 14-3-3 protein and suppression of cell death (6). Oxidative stress induces dephosphorylation of Ser967 and phosphorylation of Thr845 in the activation loop of ASK1, and both are correlated with ASK1 activity and ASK1-dependent apoptosis (7,8). On the other hand, Akt phosphorylates ASK1 at Ser83, which attenuates ASK1 activity and promotes cell survival (9).

Specificity/Sensitivity: Phospho-ASK1 (Thr845) Antibody detects transfected ASK1 only when phosphorylated at threonine 845.


Source/Purification: Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide (KLH-coupled) corresponding to residues surrounding Thr845 of mouse ASK1. Antibodies are purified by protein A and peptide affinity chromatography.

Background References:

- (1) Ichijo, H. et al. (1997) *Science* 275, 90–94.
- (2) Wang, X.S. et al. (1996) *J. Biol. Chem.* 271, 31607–31611.
- (3) Matsuzawa, A. and Ichijo, H. (2001) *J. Biochem. (Tokyo)* 130, 1–8.
- (4) Tobiume, K. et al. (2001) *EMBO Rep.* 2, 222–228.
- (5) Nishitoh, H. et al. (2002) *Genes Dev.* 16, 1345–1355.
- (6) Zhang, L. et al. (1999) *Proc. Natl. Acad. Sci. USA* 96, 8511–8515.
- (7) Tobiume, K. et al. (2002) *J. Cell. Physiol.* 191, 95–104.
- (8) Goldman, E.H. et al. (2004) *J. Biol. Chem.* in press, .
- (9) Kim, A.H. et al. (2001) *Mol. Cell. Biol.* 21, 893–901.



Western blot analysis of extracts from HEK293 cells, untransfected (lane 1) or transfected with wild-type ASK1 (lane 2) or T845A mutant ASK1 (lane 3), using Phospho-ASK1 (Thr845) Antibody (upper) or ASK1 Antibody #3762 (lower). (Lysates were provided by Dr. Wang Min, Dept. of Pathology, Yale University, New Haven, CT).



Western blot analysis of extracts from COS cells, untransfected (lane 1), or transfected with wild-type ASK1 (lanes 2,3), using Phospho-ASK1 (Thr845) Antibody (upper) or ASK1 Antibody #3762 (lower). In lane 3, the lysate was treated with λ phosphatase to demonstrate phospho-specificity of Phospho-ASK1 (Thr845) Antibody.

Entrez-Gene ID # 4217
Swiss-Prot Acc. # Q99683

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at **-20°C**.
Do not aliquot the antibody.

***Species cross-reactivity is determined by western blot.**

****Anti-rabbit secondary antibodies must be used to detect this antibody.**

Recommended Antibody Dilutions:
 Western Blotting 1:1000

For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
 Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.