

# PARN (P620) Antibody

✓ 100 µl  
(10 western blots)

**Orders** ■ 877-616-CELL (2355)  
orders@cellsignaling.com

**Support** ■ 877-678-TECH (8324)  
info@cellsignaling.com

**Web** ■ www.cellsignaling.com

rev. 03/19/10

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Applications W Endogenous	Species Cross-Reactivity* H, (B, Dg)	Molecular Wt. 78 kDa	Source Rabbit**
---------------------------------	---	-------------------------	--------------------


**Background:** Cellular levels of mRNAs are controlled by mRNA stability, the rate of synthesis and the rate of degradation. The presence and length of the poly(A) tail has been associated with mRNA stability (1). Exonucleolytic shortening of the poly(A) tail is the process that initiates the decay of many eukaryotic mRNAs (2). Poly(A)-specific ribonuclease (PARN) is the enzyme responsible for initiation of deadenylation and exonucleolytic shortening of mRNA transcripts. Through an evolutionarily conserved mechanism, PARN also translationally silences selective mRNAs during early embryonic development (3). PARN is constitutively expressed in most mammalian tissues and plays a critical role in the post-transcriptional control of gene expression (4).

**Specificity/Sensitivity:** PARN (P620) Antibody detects endogenous levels of total PARN protein.

**Source/Purification:** Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to amino acid sequences surrounding Pro620 of human PARN. Antibodies are purified by Protein A and peptide affinity chromatography.

#### Background References:

- (1) Morales, J. et al. (1997) *J Biol Chem* 272, 6607–13.
- (2) Wilson, T. and Treisman, R. (1988) *Nature* 336, 396–9.
- (3) Sachs, A.B. et al. (1997) *Cell* 89, 831–8.
- (4) Fritz, D.T. et al. (2004) *Cell Biochem Biophys* 41, 265–78.



Western blot analysis of extracts from various cell lines using PARN (P620) Antibody.

Entrez-Gene ID #5073  
Swiss-Prot Acc. #O95453

**Storage:** Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

\*Species cross-reactivity is determined by western blot.

\*\*Anti-rabbit secondary antibodies must be used to detect this antibody.

#### Recommended Antibody Dilutions:

Western blotting 1:1000

For application specific protocols please see the web page for this product at [www.cellsignaling.com](http://www.cellsignaling.com).

Please visit [www.cellsignaling.com](http://www.cellsignaling.com) for a complete listing of recommended companion products.

**IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.**

**Applications Key:** W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide  
**Species Cross-Reactivity Key:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine  
 Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.