

#3938 Store at -20°C

FoxO3a (75D8) Rabbit mAb (Biotinylated)

✓ 100 µl (10 western blots)



Orders ■ 877-616-CELL (2355) orders@cellsignal.com
Support ■ 877-678-TECH (8324) info@cellsignal.com
Web ■ www.cellsignal.com

rev. 02/24/10

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.


Applications W Endogenous	Species Cross-Reactivity H, M, R	Molecular Wt. 82–97 kDa	Isotype Rabbit IgG
---------------------------------	-------------------------------------	----------------------------	-----------------------

Description: This Cell Signaling Technology antibody is conjugated to biotin under optimal conditions. The unconjugated FoxO3a (75D8) Rabbit mAb #2497 reacts with human, mouse and rat FoxO3a protein. CST expects that FoxO3a (75D8) Rabbit mAb (Biotinylated) will also recognize FoxO3a in these species.

Background: The Forkhead family of transcription factors is involved in tumorigenesis of rhabdomyosarcoma and acute leukemias (1–3). Within the family, three members (FoxO1, FoxO4 and FoxO3a) have sequence similarity to the nematode orthologue DAF-16, which mediates signaling via a pathway involving IGF1R, PI3K and Akt (4–6). Active forkhead members act as tumor suppressors by promoting cell cycle arrest and apoptosis. Increased expression of any FoxO member results in the activation of the cell cycle inhibitor p27Kip1. Forkhead transcription factors also play a part in TGF-β-mediated upregulation of p21CIP1, a process negatively regulated through PI3K (7). Increased proliferation results when forkhead transcription factors are inactivated through phosphorylation by Akt at Thr24, Ser256 and Ser319, which results in nuclear export and inhibition of transcription factor activity (8). Forkhead transcription factors can also be inhibited by the deacetylase sirtuin (SirT1) (9).

Specificity/Sensitivity: FoxO3a (75D8) Rabbit mAb (Biotinylated) detects endogenous levels of total FoxO3a protein. The antibody does not cross-react with other members of the FoxO4 or FoxO1 families.

Source/Purification: Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Glu50 of human FoxO3a.



Western blot analysis of extracts from SH-SY5 and HeLa cells using FoxO3a (75D8) Rabbit mAb (Biotinylated).

Entrez-Gene ID #2309
Swiss-Prot Acc. #O43524

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 2 mg/ml BSA, 50% glycerol. Store at -20°C. Do not aliquot antibody.

Biotinylated antibodies are designed to be detected using streptavidin conjugates.

Recommended Antibody Dilutions:
Western blotting 1:1000

For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

- Background References:**
- (1) Anderson, M.J. et al. (1998) *Genomics* 47, 187–199.
 - (2) Galili, N. et al. (1993) *Nat. Genet.* 5, 230–235.
 - (3) Borkhardt, A. et al. (1997) *Oncogene* 14, 195–202.
 - (4) Nakae, J. et al. (1999) *J. Biol. Chem.* 274, 15982–15985.
 - (5) Rena, G. et al. (1999) *J. Biol. Chem.* 274, 17179–17183.
 - (6) Guo, S. et al. (1999) *J. Biol. Chem.* 274, 17184–17192.
 - (7) Seoane, J. et al. (2004) *Cell* 117, 211–223.
 - (8) Arden, K.C. (2004) *Mol. Cell* 14, 416–418.
 - (9) Yang, Y. et al. (2005) *EMBO J.* 24, 1021–1032.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

© 2010 Cell Signaling Technology, Inc. Rabbit Monoclonal is produced under license (granting certain rights, including those under U.S. Patent Nos. 5,675,063 and 7,429,487) from Epitomics, Inc.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.