

#4485 Store at -20°C

# PTP $\mu$ (BK2) Mouse mAb



100  $\mu$ l  
 (10 Western mini-blots)

**Orders** ■ 877-616-CELL (2355)  
 orders@cellsignal.com  
**Support** ■ 877-678-TECH (8324)  
 info@cellsignal.com  
**Web** ■ www.cellsignal.com

rev. 05/02/08

Products are sold and licensed for use by end-users only for research or laboratory use. Products are not to be used on humans for clinical diagnostic, drug screening or testing purposes.

Applications	Species Cross-Reactivity*	Molecular Wt.	Source	Isotype
W, IP Endogenous	H, R, Mi, (M)	100, 110, 210 kDa	Mouse	IgG2a


**Background:** Receptor tyrosine phosphatase PTP $\mu$  has an extracellular segment characteristic of adhesion molecules: an MEM domain, an Ig domain and four fibronectin III like (FN III) repeats (1,2). PTP $\mu$  is proteolytically cleaved into two noncovalently associated fragments: one is the extracellular domain, the other includes the transmembrane and the intracellular catalytic domains. Both fragments are approximately 100 kDa (3). The extracellular domain mediates cell-cell adhesion in a homophilic, Ca<sup>2+</sup> independent manner (1,2). PTP $\mu$  associates with multiple cadherins (4). It is able to restore E-cadherin-dependent adhesion in human prostate cancer, and is required for N-cadherin-mediated neurite outgrowth (5,6). The phosphatase activity seems to be essential for the latter function but is dispensable for the former (5,6). PTP $\mu$  also associates with and recruits a scaffold protein, RACK (receptor for activated protein C kinase), to cell-cell contact sites (7). Both PKC $\delta$  and src seem to be involved in this process (6,7).

**Specificity/Sensitivity:** PTP $\mu$  Mouse mAb detects endogenous levels of total PTP $\mu$  protein. This antibody does not cross react with other receptor tyrosine phosphatases.

**Source/Purification:** Monoclonal antibody (isotype: IgG2a) is produced by immunizing mice with a synthetic peptide (KLH-coupled) corresponding to the amino-terminal residues of human PTP $\mu$ .

**Selected Application References:**

Brady-Kalnay, S.M. et al. (1995) Receptor protein tyrosine phosphatase PTP $\mu$  associates with cadherins and catenins *in vivo*. *J. Cell Biol.* 130, 977–986. Applications: IP, W.



Western blot analysis of extract from MvLu cells using PTP $\mu$  (BK2) Mouse mAb.

**Background References:**

- (1) Gebbink, M.F. et al. (1993) *J. Biol. Chem.* 268, 16101–16104.
- (2) Brady-Kalnay, S.M. and Tonks, N.K. (1994) *J. Biol. Chem.* 269, 28472–28477.
- (3) Brady-Kalnay, S.M. et al. (1998) *J. Cell Biol.* 141, 287–296.
- (4) Hellberg, C.B. et al. (2002) *J. Biol. Chem.* 277, 11165–11173.
- (5) Burden-Gulley, S.M. and Brady-Kalnay, S.M. (1999) *J. Cell Biol.* 144, 1323–1336.
- (6) Mourton, T. et al. (2001) *J. Biol. Chem.* 276, 14896–14901.

**Entrez-Gene ID** #5797  
**Swiss-Prot Acc.** #P28827

**Storage:** Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100  $\mu$ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.

**\*Species cross-reactivity is determined by Western blot.**

**Recommended Antibody Dilutions:**

Western Blotting 1:1000  
 Immunoprecipitation 1:50

**Companion Products:**

- 20X LumiGLO<sup>®</sup> Reagent and 20X Peroxide #7003
- Phospho- $\beta$ -Catenin (Ser33/37/Thr41) Antibody #9561
- $\beta$ -Catenin Antibody #9562
- Phospho- $\beta$ -Catenin (Ser45) Antibody #9564
- Phospho- $\beta$ -Catenin (Thr41/Ser45) Antibody #9565
- Phospho-PKC (pan) (BII Ser660) Antibody #9371
- Phospho-PKC Antibody Sampler Kit #9921
- Phospho-PKC $\delta$  (Thr505) Antibody #9374
- Phospho-PKC $\alpha$ / $\beta$  II (Thr638/641) Antibody #9375
- Phospho-PKC $\theta$  (Thr538) Antibody #9377
- Phospho-PKC $\zeta$ / $\lambda$  (Thr410/403) Antibody #9378
- Phospho-PKC $\delta$  (Tyr311) Antibody #2055
- Phototope<sup>®</sup>-HRP Western Blot Detection System, Anti-mouse IgG, HRP-linked Antibody #7072
- Anti-mouse IgG, HRP-linked Antibody #7076
- Prestained Protein Marker, Broad Range (Premixed Format) #7720

**IMPORTANT: For Western blots, incubate membrane with diluted antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.**

**Applications Key:** W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E—ELISA  
**Species Cross-Reactivity Key:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken X—Xenopus Z—zebra fish B—bovine Dg—Dog All—all species expected  
 Species enclosed in parentheses are predicted to react based on 100% sequence homology.

## Western Immunoblotting Protocol (Primary Antibody Incubation in Milk)

For Western blots, incubate membrane with diluted antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

### A Solutions and Reagents

**NOTE:** Prepare solutions with Milli-Q or equivalently purified water.

- 1X Phosphate Buffered Saline (PBS)
- 1X SDS Sample Buffer:** 62.5 mM Tris-HCl (pH 6.8 at 25°C), 2% w/v SDS, 10% glycerol, 50 mM DTT, 0.01% w/v bromophenol blue or phenol red
- Transfer Buffer:** 25 mM Tris base, 0.2 M glycine, 20% methanol (pH 8.5)
- 10X Tris Buffered Saline (TBS):** To prepare 1 liter of 10X TBS: 24.2 g Tris base, 80 g NaCl; adjust pH to 7.6 with HCl (use at 1X).
- Nonfat Dry Milk (weight to volume [w/v])
- Blocking Buffer:** 1X TBS, 0.1% Tween-20 with 5% w/v nonfat dry milk; for 150 ml, add 15 ml 10X TBS to 135 ml water, mix. Add 7.5 g nonfat dry milk and mix well. While stirring, add 0.15 ml Tween-20 (100%).
- Wash Buffer:** 1X TBS, 0.1% Tween-20 (TBS/T)
- Primary Antibody Dilution Buffer:** 1X TBS, 0.1% Tween-20 with 5% nonfat dry milk; for 20 ml, add 2 ml 10X TBS to 18 ml water, mix. Add 1.0 g nonfat dry milk and mix well. While stirring, add 20 µl Tween-20 (100%).
- Phototope®-HRP Western Blot Detection System #7072:** Includes biotinylated protein ladder, secondary anti-mouse (#7076) antibody conjugated to horseradish peroxidase (HRP), anti-biotin antibody conjugated to HRP, LumiGLO® chemiluminescent reagent and peroxide.
- Prestained Protein Marker, Broad Range (Premixed Format) #7720
- Biotinylated Protein Ladder Detection Pack #7727
- Blotting Membrane:** This protocol has been optimized for nitrocellulose membranes, which CST recommends. PVDF membranes may also be used.

### B Protein Blotting

A general protocol for sample preparation is described below.

- Treat cells by adding fresh media containing regulator for desired time.
- Aspirate media from cultures; wash cells with 1X PBS; aspirate.
- Lyse cells by adding 1X SDS sample buffer (100 µl per well of 6-well plate or 500 µl per plate of 10 cm diameter plate). Immediately scrape the cells off the plate and transfer the extract to a microcentrifuge tube. Keep on ice.
- Sonicate for 10–15 seconds to shear DNA and reduce sample viscosity.
- Heat a 20 µl sample to 95–100°C for 5 minutes; cool on ice.
- Microcentrifuge for 5 minutes.
- Load 20 µl onto SDS-PAGE gel (10 cm x 10 cm).

**NOTE:** CST recommends loading prestained molecular weight marker (#7720, 10 µl/lane) to verify electrotransfer and biotinylated protein ladder (#7727, 10 µl/lane) to determine molecular weights.

- Electrotransfer to nitrocellulose or PVDF membrane.

### C Membrane Blocking and Antibody Incubations

**NOTE:** Volumes are for 10 cm x 10 cm (100 cm<sup>2</sup>) of membrane; for different sized membranes, adjust volumes accordingly.

- (Optional) After transfer, wash nitrocellulose membrane with 25 ml TBS for 5 minutes at room temperature.
- Incubate membrane in 25 ml of blocking buffer for 1 hour at room temperature.
- Wash three times for 5 minutes each with 15 ml of TBS/T.
- Incubate membrane and primary antibody (at the appropriate dilution) in 10 ml primary antibody dilution buffer with gentle agitation overnight at 4°C.
- Wash three times for 5 minutes each with 15 ml of TBS/T.
- Incubate membrane with HRP-conjugated secondary antibody (1:2000) and HRP-conjugated anti-biotin antibody (1:1000) to detect biotinylated protein markers in 10 ml of blocking buffer with gentle agitation for 1 hour at room temperature.
- Wash three times for 5 minutes each with 15 ml of TBS/T.

### D Detection of Proteins

- Incubate membrane with 10 ml LumiGLO® (0.5 ml 20X LumiGLO®, 0.5 ml 20X Peroxide and 9.0 ml Milli-Q water) with gentle agitation for 1 minute at room temperature.

**NOTE:** LumiGLO® substrate can be further diluted if signal response is too fast.

- Drain membrane of excess developing solution (do not let dry), wrap in plastic wrap and expose to x-ray film. An initial 10-second exposure should indicate the proper exposure time.

**NOTE:** Due to the kinetics of the detection reaction, signal is most intense immediately following LumiGLO® incubation and declines over the following 2 hours.

## Immunoprecipitation Protocol / (For Analysis By Western Immunoblotting)

### A Solutions and Reagents

**NOTE:** Prepare solutions with Milli-Q or equivalently purified water.

1. 1X Phosphate Buffered Saline (PBS)
2. **1X Cell Lysis Buffer:** 20 mM Tris (pH 7.5), 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% Triton X-100, 2.5 mM Sodium pyrophosphate, 1 mM  $\beta$ -glycerophosphate, 1 mM  $\text{Na}_3\text{VO}_4$ , 1  $\mu\text{g/ml}$  Leupeptin

**NOTE:** Add 1 mM PMSF immediately prior to use.

1. **Transfer Buffer:** 25 mM Tris base, 0.2 mM glycine, 20% methanol (pH 8.5)
2. **Protein A or G Agarose Beads:** (Can be stored for 2 weeks at 4°C.) Please prepare according to manufacturer's instructions. Use Protein A for rabbit IgG pull down and Protein G for mouse IgG pull down.
3. **3X SDS Sample Buffer:** 187.5 mM Tris-HCl (pH 6.8 at 25°C), 6% w/v SDS, 30% glycerol, 150 mM DTT, 0.03% w/v bromophenol blue

### B Preparing Cell Lysates

1. Aspirate media. Treat cells by adding fresh media containing regulator for desired time.
2. To harvest cells under nondenaturing conditions, remove media and rinse cells once with ice-cold PBS.
3. Remove PBS and add 0.5 ml ice-cold 1X cell lysis buffer to each plate (10 cm) and incubate the plates on ice for 5 minutes.
4. Scrape cells off the plates and transfer to microcentrifuge tubes. Keep on ice.
5. Sonicate samples on ice three times for 5 seconds each.
6. Microcentrifuge for 10 minutes at 14,000 X g, 4°C, and transfer the supernatant to a new tube. If necessary, lysate can be stored at -80°C.

### C Immunoprecipitation

Optional: It may be necessary to perform a lysate pre-clearing step to reduce non-specific binding to the Protein A/G agarose beads (See section below).

1. Take 200  $\mu\text{l}$  cell lysate and add primary antibody. Incubate with gentle rocking overnight at 4°C.
2. Add either protein A or G agarose beads (20  $\mu\text{l}$  of 50% bead slurry). Incubate with gentle rocking for 1–3 hours at 4°C.
3. Microcentrifuge for 30 seconds at 4°C. Wash pellet five times with 500  $\mu\text{l}$  of 1X cell lysis buffer. Keep on ice during washes.
4. Resuspend the pellet with 20  $\mu\text{l}$  3X SDS sample buffer. Vortex, then microcentrifuge for 30 seconds.
5. Heat the sample to 95–100°C for 2–5 minutes and microcentrifuge for 1 minute at 14,000 X g.
6. Load the sample (15–30  $\mu\text{l}$ ) on SDS-PAGE gel (12–15%).
7. Analyze sample by Western blotting (see Western Immunoblotting Protocol).

### Cell Lysate Pre-Clearing (Optional)

1. Take 200  $\mu\text{l}$  cell lysate and add to either Protein A or G agarose beads (20  $\mu\text{l}$  of 50% bead slurry).
2. Incubate at 4°C for 30 – 60 minutes.
3. Spin for 10 minutes at 4°C. Transfer the supernatant to a fresh tube.
4. Proceed to step 1 of Immunoprecipitation.