

Phospho-IκB-β (Thr19/Ser23) Antibody (Human Specific)



Cell Signaling
TECHNOLOGY®

✓ 100 μl
(10 Western mini-blot)

Orders ■ 877-616-CELL (2355)
orders@cellsignal.com

Support ■ 877-678-TECH (8324)
info@cellsignal.com

Web ■ www.cellsignal.com

rev. 01/08/07

This product is for *in vitro* research use only and is not intended for use in humans or animals.

Applications W	Species Cross-Reactivity H	Molecular Wt. 48, 50 kDa	Source Rabbit
-------------------	-------------------------------	-----------------------------	------------------


Background: The NF-κB/Rel transcription factors are present in the cytosol in an inactive state complexed with the inhibitory IκB proteins (1–3). Activation occurs via phosphorylation of IκB-α at Ser32 and Ser36 followed by proteasome-mediated degradation, resulting in the release and nuclear translocation of active NF-κB (3–7). IκB-α phosphorylation and resulting Rel-dependent transcription are activated by a highly diverse group of extracellular signals including inflammatory cytokines, growth factors and chemokines.

Kinases that phosphorylate IκB at these activating sites have been identified (8). Because phosphorylation of IκB-α at Ser32 is essential for release of active NF-κB, phosphorylation at this site is an excellent marker of NF-κB activation (1–3).

The regulation of IκB-β and IκB-ε is similar to that of IκB-α. However, the phosphorylation and ubiquitin-mediated degradation of these proteins occurs with much slower kinetics (9). IKK phosphorylation of IκB-β occurs at Ser19 and Ser23, while IκBε can be phosphorylated at Ser18 and Ser22 (10). The human sequence of IκB-β has also been reported to contain a threonine at position 19 suggesting that phosphorylation could be Thr19/Ser23 (11).

Specificity/Sensitivity: Phospho-IκB-β (Thr19/Ser23) Antibody detects endogenous levels of human IκB-β only when phosphorylated at threonine 19 and serine 23. This antibody also recognizes phosphorylation at Ser19/Ser23 also reported as the sequence for IκB-β.

Source/Purification: Polyclonal antibodies are produced by immunizing rabbits with a synthetic phosphopeptide (KLH-coupled) corresponding to residues of human IκB-β surrounding Thr19/Ser23. Antibodies are purified by protein A and affinity chromatography.




Western blot analysis of extracts from HT-29 cells treated with TNF-α (20 ng/ml) and calyculin A (100 nM) for 10 minutes as indicated, using Phospho-IκB-β (Thr19/Ser23) Antibody (upper) or IκB-β Antibody #9248 (lower).

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Recommended Antibody Dilutions:
Western blotting 1:1000

- Companion Products:**
- IκB-α Antibody #9242
 - IκB-α (112B2) Mouse mAb (Carboxy-Terminal Antigen) #9247
 - IκB-β Antibody #9248
 - IκB-ε Antibody #9249
 - Phospho-IκB-α (Ser32) Antibody #9241
 - Phospho-IκB-α (Ser32/36) (12C2) Monoclonal Antibody #5210
 - Phospho-IκB-α (Ser32/36) (5A5) Monoclonal Antibody #5205
 - Phospho-IκB-α (Ser32/36) (5A5) Mouse mAb #9246
 - Phospho-IκBβ (Ser19/23) Antibody (Mouse/Rat Specific) #9245
 - Phospho-IκB-ε (Ser18/22) Antibody #4924
 - Phototope®-HRP Western Detection System, Anti-rabbit IgG, HRP-linked Antibody #7071
 - Anti-rabbit IgG, HRP-linked Antibody #7074
 - Prestained Protein Marker, Broad Range (Premixed Format) #7720

Biotinylated Protein Ladder Detection Pack #7727
LumiGLO® Reagent and Peroxide #7003



NFκB Signaling Pathway

IMPORTANT: For Western blots, incubate membrane with diluted antibody in 5% BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry IC—Immunocytochemistry IF—Immunofluorescence F—Flow cytometry E—ELISA D—DELIA®
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken X—Xenopus Z—zebra fish B—bovine All—all species expected
Species enclosed in parentheses are predicted to react based on 100% sequence homology.

Background References:

- (1) Baeuerle, P.A. and Baltimore, D. (1988) *Science* 242, 540–546.
- (2) Beg, A.A. et al. (1993) *Genes Dev.* 7, 2064–2070.
- (3) Finco, T.S. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91, 11884–11888.
- (4) Brown, K. et al. (1995) *Science* 267, 1485–1488.
- (5) Brockman, J.A. et al. (1995) *Mol. Cell. Biol.* 15, 2809–2818.
- (6) Traenckner, E.B. et al. (1995) *EMBO J.* 14, 2876–2883.
- (7) Chen, Z.J. et al. (1996) *Cell* 84, 853–862.
- (8) Karin, M. and Ben-Neriah, Y. (2000) *Annu. Rev. Immunol.* 18, 621–663.
- (9) Hoffmann, A. et al. (2002) *Science* 298, 1241–1245.
- (10) Shirane, M. et al. (1999) *J. Biol. Chem.* 274, 28109–28174.
- (11) Lee, J.W. et al. (1995) *Mol. Endocrinol.* 9, 243–254.

For Western blots, incubate membrane with diluted anti-body in 5% BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

A. Solutions and Reagents

Note: Prepare solutions with Milli-Q or equivalently purified water.

- A1. ■ 1X Phosphate Buffered Saline (PBS)
- A2. ■ 1X SDS Sample Buffer:
62.5 mM Tris-HCl (pH 6.8 at 25°C), 2% w/v SDS, 10% glycerol, 50 mM DTT, 0.01% w/v bromophenol blue or phenol red
- A3. ■ Transfer Buffer:
25 mM Tris base, 0.2 M glycine, 20% methanol (pH 8.5)
- A4. ■ 10X Tris Buffered Saline (TBS):
To prepare 1 liter of 10X TBS: 24.2 g Tris base, 80 g NaCl; adjust pH to 7.6 with HCl (use at 1X).
- A5. ■ Nonfat Dry Milk (weight to volume [w/v])
- A6. ■ Blocking Buffer:
1X TBS, 0.1% Tween-20 with 5% w/v nonfat dry milk; for 150 ml, add 15 ml 10X TBS to 135 ml water, mix. Add 7.5 g nonfat dry milk and mix well. While stirring, add 0.15 ml Tween-20 (100%).
- A7. ■ Wash Buffer:
1X TBS, 0.1% Tween-20 (TBS/T)
- A8. ■ Bovine Serum Albumin (BSA)
- A9. ■ Primary Antibody Dilution Buffer:
1X TBS, 0.1% Tween-20 with 5% BSA; for 20 ml, add 2 ml 10X TBS to 18 ml water, mix. Add 1.0 g BSA and mix well. While stirring, add 20 µl Tween-20 (100%).
- A10. ■ Phototope®-HRP Western Blot Detection System #7071:
Includes biotinylated protein ladder, secondary anti-rabbit (#7074) antibody conjugated to horseradish peroxidase (HRP), anti-biotin antibody conjugated to HRP, LumiGLO® chemiluminescent reagent and peroxide.
- A11. ■ Prestained Protein Marker, Broad Range (Premixed Format) #7720
- A12. ■ Biotinylated Protein Ladder Detection Pack #7727
- A13. ■ Blotting Membrane:
This protocol has been optimized for nitrocellulose membranes, which CST recommends. PVDF membranes may also be used.

B. Protein Blotting

A general protocol for sample preparation is described below.

- B1. Treat cells by adding fresh media containing regulator for desired time.
- B2. Aspirate media from cultures; wash cells with 1X PBS; aspirate.
- B3. Lyse cells by adding 1X SDS sample buffer (100 µl per well of 6-well plate or 500 µl per plate of 10 cm diameter plate). Immediately scrape the cells off the plate and transfer the extract to a microcentrifuge tube. Keep on ice.
- B4. Sonicate for 10–15 seconds to shear DNA and reduce sample viscosity.
- B5. Heat a 20 µl sample to 95–100°C for 5 minutes; cool on ice.
- B6. Microcentrifuge for 5 minutes.
- B7. Load 20 µl onto SDS-PAGE gel (10 cm x 10 cm).

Note: CST recommends loading prestained molecular weight markers (#7720, 10 µl/lane) to verify electrotransfer and biotinylated protein ladder (#7727, 10 µl/lane) to determine molecular weights.

- B8. Electrotransfer to nitrocellulose or PVDF membrane.

C. Membrane Blocking and Antibody Incubations

Note: Volumes are for 10 cm x 10 cm (100 cm²) of membrane; for different sized membranes, adjust volumes accordingly.

- C1. (Optional) After transfer, wash nitrocellulose membrane with 25 ml TBS for 5 minutes at room temperature.
- C2. Incubate membrane in 25 ml of blocking buffer for 1 hour at room temperature.
- C3. Wash three times for 5 minutes each with 15 ml of TBS/T.
- C4. Incubate membrane and primary antibody (at the appropriate dilution) in 10 ml primary antibody dilution buffer with gentle agitation overnight at 4°C.
- C5. Wash three times for 5 minutes each with 15 ml of TBS/T.
- C6. Incubate membrane with HRP-conjugated secondary antibody (1:2000) and HRP-conjugated anti-biotin antibody (1:1000) to detect biotinylated protein markers in 10 ml of blocking buffer with gentle agitation for 1 hour at room temperature.
- C7. Wash three times for 5 minutes each with 15 ml of TBS/T.

D. Detection of Proteins

- D1. Incubate membrane with 10 ml LumiGLO® (0.5 ml 20X LumiGLO®, 0.5 ml 20X Peroxide and 9.0 ml Milli-Q water) with gentle agitation for 1 minute at room temperature.

Note: LumiGLO® substrate can be further diluted if signal response is too fast.

- D2. Drain membrane of excess developing solution (do not let dry), wrap in plastic wrap and expose to x-ray film. An initial 10-second exposure should indicate the proper exposure time.

Note: Due to the kinetics of the detection reaction, signal is most intense immediately following LumiGLO® incubation and declines over the following 2 hours.