

# Phospho-Elk-1 (Ser383) (2B1) Mouse mAb

✓ 100 µl  
(10 western blots)

Orders ■ 877-616-CELL (2355)  
orders@cellsignal.com  
Support ■ 877-678-TECH (8324)  
info@cellsignal.com  
Web ■ www.cellsignal.com

rev. 12/16/09

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Entrez-Gene ID #2002  
Swiss-Prot Acc. #P19419

Applications W Recombinant	Species Cross-Reactivity* (H, M, R)	Molecular Wt. 47 kDa	Isotype Mouse IgG1**
----------------------------------	--	-------------------------	-------------------------


**Background:** The transcription factor Elk-1 is a component of the ternary complex that binds the serum response element (SRE) and mediates gene activity in response to serum and growth factors (1–3). Elk-1 is phosphorylated by MAP kinase pathways at a cluster of S/T motifs at its carboxy-terminus; phosphorylation at these sites, particularly Ser383, is critical for transcriptional activation by Elk-1. Elk-1 appears to be a direct target of activated MAP kinase: (a) biochemical studies indicate that Elk-1 is a good substrate for MAP kinase; (b) the kinetics of Elk-1 phosphorylation and activation correlate with MAP kinase activity; (c) interfering mutants of MAP kinase block Elk-1 activation *in vivo*. Other studies have shown that Elk-1 (Ser383) is also a target of the stress-activated kinase SAPK/JNK (4,5).

**Specificity/Sensitivity:** Phospho-Elk-1 (Ser383) (2B1) Mouse mAb detects less than 5 ng of phosphorylated Elk-1, and will not react with up to 1 µg of nonphosphorylated Elk-1 protein.

**Source/Purification:** Monoclonal antibody is produced by immunizing animals with a synthetic phospho-peptide (KLH-coupled) corresponding to residues around Ser383 of human Elk-1.

**Background References:**

- (1) Marais, R. et al. (1993) *Cell* 73, 381–393.
- (2) Kortenjann, M. et al. (1994) *Mol. Cell. Biol.* 14, 4815–4824.
- (3) Hill, C.S. and Treisman, R. (1995) *Cell* 80, 199–211.
- (4) Cavigelli, M. et al. (1995) *EMBO J.* 14, 5957–5964.
- (5) Whitmarsh, A.J. et al. (1995) *Science* 269, 403–407.



Western blot analysis of Elk-1 fusion protein expressed from *E. coli* with or without phosphorylation with purified Erk2 enzyme, using Phospho-Elk-1 (Ser383) (2B1) Mouse mAb (upper) or Elk-1 Antibody #9182 (lower).

**Storage:** Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

\*Species cross-reactivity is determined by western blot.

\*\*Anti-mouse secondary antibodies must be used to detect this antibody.

**Recommended Antibody Dilutions:**

Western blotting 1:1000

For application specific protocols please see the web page for this product at [www.cellsignal.com](http://www.cellsignal.com).

Please visit [www.cellsignal.com](http://www.cellsignal.com) for a complete listing of recommended companion products.

**IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.**

**Applications Key:** W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide  
**Species Cross-Reactivity Key:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine  
Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.