

Bad (D24A9) Rabbit mAb

✓ 100 µl
(10 western blots)

Orders ■ 877-616-CELL (2355)
orders@cellsignal.com

Support ■ 877-678-TECH (8324)
info@cellsignal.com

Web ■ www.cellsignal.com

rev. 02/08/10

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Entrez-Gene ID #572
Swiss-Prot Acc. #Q92934

Applications	Species Cross-Reactivity*	Molecular Wt.	Isotype
W Endogenous	H, M, R, Mk	23 kDa	Rabbit IgG**


Background: Bad is a proapoptotic member of the Bcl-2 family that promotes cell death by displacing Bax from binding to Bcl-2 and Bcl-xL (1,2). Survival factors, such as IL-3, inhibit the apoptotic activity of Bad by activating intracellular signaling pathways that result in the phosphorylation of Bad at Ser112 and Ser136 (2). Phosphorylation at these sites promotes binding of Bad to 14-3-3 proteins to prevent an association between Bad with Bcl-2 and Bcl-xL (2). Akt phosphorylates Bad at Ser136 to promote cell survival (3,4). Bad is phosphorylated at Ser112 both *in vivo* and *in vitro* by p90RSK (5,6) and mitochondria-anchored PKA (7). Phosphorylation of Ser155 in the BH3 domain by PKA plays a critical role in blocking the dimerization of Bad and Bcl-xL (8-10).

Specificity/Sensitivity: Bad (D24A9) Rabbit mAb detects endogenous levels of total Bad protein. The antibody does not cross-react with related proteins.

Source/Purification: Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro102 of human Bad.

Background References:

- (1) Yang, E. et al. (1995) *Cell* 80, 285-291.
- (2) Zha, J. et al. (1996) *Cell* 87, 619-628.
- (3) Datta, S.R. et al. (1997) *Cell* 91, 231-241.
- (4) Peso, L. et al. (1997) *Science* 278, 687-689.
- (5) Bonni, A. et al. (1999) *Science* 286, 1358-1362.
- (6) Tan, Y. et al. (1999) *J. Biol. Chem.* 274, 34859-34867.
- (7) Harada, H. et al. (1999) *Mol. Cell* 3, 413-422.
- (8) Tan, Y. et al. (2000) *J. Biol. Chem.* 275, 25865-25869.
- (9) Lizcano, J. et al. (2000) *Biochem. J.* 349, 547-557.
- (10) Datta, S. et al. (2000) *Mol. Cell* 6, 41-51.



Western blot analysis of extracts from various cell lines using Bad (D24A9) Rabbit mAb.

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

*Species cross-reactivity is determined by western blot.

**Anti-rabbit secondary antibodies must be used to detect this antibody.

Recommended Antibody Dilutions:

Western blotting 1:1000

For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide

Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebra fish B—bovine

Dg—dog Pg—pig Sc—S. cerevisiae All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.