

#9320 Store at -20°C

Src Family Antibody Sampler Kit



1 Kit
(6 x 40 µl)

Orders ■ 877-616-CELL (2355)
orders@cellsignal.com
Support ■ 877-678-TECH (8324)
info@cellsignal.com
Web ■ www.cellsignal.com

New 12/09

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Products Included	Product #	Quantity	Mol. Wt.	Isotype
Src (32G6) Rabbit mAb	2123	40 µl	60 kDa	Rabbit IgG
Csk (C74C1) Rabbit mAb	4980	40 µl	50 kDa	Rabbit IgG
Fyn Antibody	4023	40 µl	59 kDa	Rabbit IgG
Lck (D88) XP™ Rabbit mAb	2984	40 µl	56 kDa	Rabbit IgG
Lyn (C13F9) Rabbit mAb	2796	40 µl	56 kDa	Rabbit IgG
Yes Antibody	2734	40 µl	60 kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

See www.cellsignal.com for individual component applications, species cross-reactivity, dilutions and additional application protocols.

Description: The Src Family Antibody Sampler Kit provides an economical means of evaluating total levels of Src family member proteins. The kit contains enough primary and secondary antibody to perform four western blots with each antibody.

Background: The Src family of protein tyrosine kinases, which includes Src, Lyn, Fyn, Yes, Lck, Blk and Hck, are important in the regulation of growth and differentiation of eukaryotic cells (1). Src activity is regulated by tyrosine phosphorylation at two sites, but with opposing effects. Phosphorylation of Src at Tyr416 in the activation loop of the kinase domain by carboxy-terminal Src kinase (Csk) upregulates enzyme activity, whereas phosphorylation of Tyr527 in the carboxy-terminal tail renders the enzyme less active (2).

The kinase Fyn plays a role in T-cell receptor signaling and also helps in memory formation and adhesion mediated signaling (3,4). Lymphocyte specific kinase Lck is essential in the differentiation and activation of T-cells (3). Hematopoietic Lyn kinase, also phosphorylated by Csk, is involved in the regulation of B-cell function, migration and development (5). The ubiquitously expressed kinase Yes acts downstream of several different cell surface receptors, including G-protein-coupled receptors, and is involved in the regulation of angiogenesis, the cell cycle and cell adhesion (6,7).

Specificity/Sensitivity: Each antibody in the Src Family Antibody Sampler Kit detects endogenous levels of its target protein and does not cross-react with other family members.

Source/Purification: Src (32G6) Rabbit mAb is produced by immunizing animals with a Src fusion protein containing amino acid residues 1-110 of human Src. The remaining monoclonal antibodies are produced by immunizing animals with synthetic peptides (KLH-coupled) corresponding to residues surrounding Val399 of human Csk, Tyr505 of human Lck, and the amino-terminal sequence of human Lyn. Polyclonal antibodies are produced by immunizing animals with a synthetic peptide (KLH-coupled) corresponding to residues surrounding Ser25 of human Fyn and the amino terminus of human Yes. Polyclonal antibodies are purified by protein A and peptide affinity chromatography.


Background References:

- (1) Thomas, S.M. and Brugge, J.S. (1997) *Annu. Rev. Cell Dev. Biol.* 13, 513-609.
- (2) Hunter, T. (1987) *Cell* 49, 1-4.
- (3) Palacios, E.H. and Weiss, A. (2004) *Oncogene* 23, 7990-8000.
- (4) Isosaka, T. et al. (2008) *Eur. J. Neurosci.* 28, 973-981.
- (5) Seo, S. et al. (2001) *J. Immunol.* 166, 3710-3716.
- (6) Summy, J.M. et al. (2003) *J. Cell Sci.* 116, 2585-2598.
- (7) Werdich, X.Q. and Penn, J.S. (2005) *Angiogenesis* 8, 315-326.


Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibodies.

Recommended Antibody Dilutions:
Western blotting 1:1000


Please visit www.cellsignal.com for a complete listing of recommended companion products.




Western blot analysis of extracts from various cell lines using **Csk (C74C1) Rabbit mAb #4980**.




Western blot analysis of extracts from various cell lines using **Fyn Antibody #4023**.




Western blot analysis of cell extracts from various cell lines using **Src (32G6) Rabbit mAb #2123**.



Western blot analysis of extracts from various cell lines using **Lyn (C13F9) Rabbit mAb #2796**.



Western blot analysis of extracts from Ramos and Jurkat cells using **Lck (D88) XP™ Rabbit mAb #2984**.



Western blot analysis of lysates from SEM and U-87 MG cells using **Yes Antibody #2734**.

Western Immunoblotting Protocol (Primary Antibody Incubation in BSA)

For Western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

A Solutions and Reagents

NOTE: Prepare solutions with Milli-Q or equivalently purified water.

- 1X Phosphate Buffered Saline (PBS)
- 1X SDS Sample Buffer:** 62.5 mM Tris-HCl (pH 6.8 at 25°C), 2% w/v SDS, 10% glycerol, 50 mM DTT, 0.01% w/v bromophenol blue or phenol red
- Transfer Buffer:** 25 mM Tris base, 0.2 M glycine, 20% methanol (pH 8.5)
- 10X Tris Buffered Saline (TBS):** To prepare 1 liter of 10X TBS: 24.2 g Tris base, 80 g NaCl; adjust pH to 7.6 with HCl (use at 1X).
- Nonfat Dry Milk (weight to volume [w/v])
- Blocking Buffer:** 1X TBS, 0.1% Tween-20 with 5% w/v nonfat dry milk; for 150 ml, add 15 ml 10X TBS to 135 ml water, mix. Add 7.5 g nonfat dry milk and mix well. While stirring, add 0.15 ml Tween-20 (100%).
- Wash Buffer:** 1X TBS, 0.1% Tween-20 (TBS/T)
- Bovine Serum Albumin (BSA)
- Primary Antibody Dilution Buffer:** 1X TBS, 0.1% Tween-20 with 5% BSA; for 20 ml, add 2 ml 10X TBS to 18 ml water, mix. Add 1.0 g BSA and mix well. While stirring, add 20 µl Tween-20 (100%).
- Phototope[®]-HRP Western Blot Detection System #7071:** Includes biotinylated protein ladder, secondary anti-rabbit (#7074) antibody conjugated to horseradish peroxidase (HRP), anti-biotin antibody conjugated to HRP, LumiGLO[®] chemiluminescent reagent and peroxide.
- Prestained Protein Marker, Broad Range (Premixed Format) #7720
- Biotinylated Protein Ladder Detection Pack #7727
- Blotting Membrane:** This protocol has been optimized for nitrocellulose membranes, which CST recommends. PVDF membranes may also be used.

B Protein Blotting

A general protocol for sample preparation is described below.

- Treat cells by adding fresh media containing regulator for desired time.
- Aspirate media from cultures; wash cells with 1X PBS; aspirate.
- Lyse cells by adding 1X SDS sample buffer (100 µl per well of 6-well plate or 500 µl per plate of 10 cm diameter plate). Immediately scrape the cells off the plate and transfer the extract to a microcentrifuge tube. Keep on ice.
- Sonicate for 10–15 seconds to shear DNA and reduce sample viscosity.
- Heat a 20 µl sample to 95–100°C for 5 minutes; cool on ice.
- Microcentrifuge for 5 minutes.
- Load 20 µl onto SDS-PAGE gel (10 cm x 10 cm).

NOTE: CST recommends loading prestained molecular weight markers (#7720, 10 µl/lane) to verify electrotransfer and biotinylated protein ladder (#7727, 10 µl/lane) to determine molecular weights.

- Electrotransfer to nitrocellulose or PVDF membrane.

C Membrane Blocking and Antibody Incubations

NOTE: Volumes are for 10 cm x 10 cm (100 cm²) of membrane; for different sized membranes, adjust volumes accordingly.

- (Optional) After transfer, wash nitrocellulose membrane with 25 ml TBS for 5 minutes at room temperature.
- Incubate membrane in 25 ml of blocking buffer for 1 hour at room temperature.
- Wash three times for 5 minutes each with 15 ml of TBS/T.
- Incubate membrane and primary antibody (at the appropriate dilution) in 10 ml primary antibody dilution buffer with gentle agitation overnight at 4°C.
- Wash three times for 5 minutes each with 15 ml of TBS/T.
- Incubate membrane with HRP-conjugated secondary antibody (1:2000) and HRP-conjugated anti-biotin antibody (1:1000) to detect biotinylated protein markers in 10 ml of blocking buffer with gentle agitation for 1 hour at room temperature.
- Wash three times for 5 minutes each with 15 ml of TBS/T.

D Detection of Proteins

- Incubate membrane with 10 ml LumiGLO[®] (0.5 ml 20X LumiGLO[®], 0.5 ml 20X Peroxide and 9.0 ml Milli-Q water) with gentle agitation for 1 minute at room temperature.

NOTE: LumiGLO[®] substrate can be further diluted if signal response is too fast.

- Drain membrane of excess developing solution (do not let dry), wrap in plastic wrap and expose to x-ray film. An initial 10-second exposure should indicate the proper exposure time.

NOTE: Due to the kinetics of the detection reaction, signal is most intense immediately following LumiGLO[®] incubation and declines over the following 2 hours.