

#9368 Store at -20°C

PKCζ (C24E6) Rabbit mAb

✓ 100 µl
(10 Western mini-blot)



Orders ■ 877-616-CELL (2355)
orders@cellsignal.com
Support ■ 877-678-TECH (8324)
info@cellsignal.com
Web ■ www.cellsignal.com

New 12/06

This product is for *in vitro* research use only and is not intended for use in humans or animals.

| Applications | Species Cross-Reactivity | Molecular Wt. | Source | Isotype |
|--------------|--------------------------|---------------|---------|---------|
| W | H, M, R, Mk | 78 kDa | Rabbit* | IgG |


Background: Activation of PKC is one of the earliest events in a cascade leading to a variety of cellular responses such as secretion, gene expression, proliferation and muscle contraction (1,2). PKC isoforms have been classified into three groups: classical PKCs, which are Ca²⁺ dependent via their C2 domains and are activated by phosphatidylserine (PS), diacylglycerol (DAG) and phorbol esters (TPA or PMA) through their cysteine-rich C1 domains, novel PKCs, which are Ca²⁺ independent but are still regulated by PS, DAG and TPA and atypical PKCs, which are Ca²⁺ independent and do not require PS, DAG or TPA for their activation (3–7). Members of these three PKC groups contain a pseudo-substrate or autoinhibitory domain that binds to the substrate binding site in the catalytic domain, preventing its activation in the absence of cofactors or activators.

Other members have been recently added to the PKC superfamily based on homology within the catalytic domain. PKC, or PKD, is regulated by DAG and TPA through its C1 domain. However, PKD is distinguished by a PH domain, as well as by its unique substrate recognition and Golgi localization. The PKC-related kinases, or PRKs, lack a C1 domain and thus do not respond to DAG or phorbol esters. Instead, they can be activated by phosphatidylinositol lipids and their kinase activity is directly regulated by small GTPases of the Rho family through Rho binding to the homology region 1 (HR1).

The activity of PKC is under the control of three distinct phosphorylation events. Specifically, Thr500 in the activation loop, the Thr641 autophosphorylation site and the Ser660 hydrophobic site at the carboxy terminus of PKCβII are phosphorylated *in vivo* (2). For the atypical PKC isoforms, there is no phosphorylation in the hydrophobic region, which has a glutamic acid residue in place of the serine or threonine residue found in other PKC isoforms. The enzyme PDK1, or perhaps a close relative, is responsible for PKC activation.

Specificity/Sensitivity: PKCζ (C24E6) Rabbit mAb detects endogenous levels of total PKCζ. The antibody does not cross-react with other PKC isoforms.

Source/Purification: Monoclonal antibodies are produced by immunizing rabbits with a synthetic peptide (KLH-coupled) corresponding to human PKCζ.



Western blot analysis of extracts from 293, NBT-II, PC12 and COS cells, using PKCζ (C24E6) Rabbit mAb.

Background References:

- (1) Nishizuka, Y. (1984) *Nature* 308, 693–698.
- (2) Keranen, L.M. et al. (1995) *Curr. Biol.* 5, 1394–1403.
- (3) Newton, A.C. (1995) *J. Biol. Chem.* 270, 28495–28498.
- (4) Mellor, H. and Parker, P.J. (1998) *Biochem. J.* 332, 281–292.
- (5) Ron, D. and Kazanietz, M.G. (1999) *FASEB J.* 13, 1658–1676.
- (6) Way, K.J. et al. (2000) *Trends Pharmacol. Sci.* 21, 181–187.
- (7) Moscat, J. and Diaz-Meco, M.T. (2000) *EMBO Rep.* 1, 399–403.
- (8) Good, J.A. et al. (1998) *Science* 281, 2042–2045.
- (9) Sonnenburg, E.D. (2001) *J. Biol. Chem.* 276, 45289–45297.

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

***Anti-rabbit secondary antibodies must be used to detect this antibody**

Recommended Antibody Dilutions:
Western blotting 1:1000

Companion Products:

- PKCζ Antibody #9372
- PKCα Antibody #2056
- PKCδ Antibody #2058
- PKCθ Antibody #2059
- PKD/PKCμ Antibody #2052
- Phototope®-HRP Western Blot Detection System, Anti-rabbit IgG, HRP-linked Antibody #7071
- Anti-rabbit IgG, HRP-linked Antibody #7074
- Prestained Protein Marker, Broad Range (Premixed Format) #7720
- Biotinylated Protein Ladder Detection Pack #7727
- 20X LumiGLO® Reagent and 20X Peroxide #7003

IMPORTANT: For Western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry IC—Immunocytochemistry IF—Immunofluorescence
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken X—Xenopus
 F—Flow cytometry E—ELISA D—DELFIATM
 Z—zebra fish B—bovine All—all species expected
 Species enclosed in parentheses are predicted to react based on 100% sequence homology.

Western Immunoblotting Protocol (Primary Ab Incubation In BSA)

For Western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

A Solutions and Reagents

NOTE: Prepare solutions with Milli-Q or equivalently purified water.

- 1X Phosphate Buffered Saline (PBS)
- 1X SDS Sample Buffer:** 62.5 mM Tris-HCl (pH 6.8 at 25°C), 2% w/v SDS, 10% glycerol, 50 mM DTT, 0.01% w/v bromophenol blue or phenol red
- Transfer Buffer:** 25 mM Tris base, 0.2 M glycine, 20% methanol (pH 8.5)
- 10X Tris Buffered Saline (TBS):** To prepare 1 liter of 10X TBS: 24.2 g Tris base, 80 g NaCl; adjust pH to 7.6 with HCl (use at 1X).
- Nonfat Dry Milk (weight to volume [w/v])
- Blocking Buffer:** 1X TBS, 0.1% Tween-20 with 5% w/v nonfat dry milk; for 150 ml, add 15 ml 10X TBS to 135 ml water, mix. Add 7.5 g nonfat dry milk and mix well. While stirring, add 0.15 ml Tween-20 (100%).
- Wash Buffer:** 1X TBS, 0.1% Tween-20 (TBS/T)
- Bovine Serum Albumin (BSA)
- Primary Antibody Dilution Buffer:** 1X TBS, 0.1% Tween-20 with 5% BSA; for 20 ml, add 2 ml 10X TBS to 18 ml water, mix. Add 1.0 g BSA and mix well. While stirring, add 20 µl Tween-20 (100%).
- Phototope[®]-HRP Western Blot Detection System #7071:** Includes biotinylated protein ladder, secondary anti-rabbit (#7074) antibody conjugated to horseradish peroxidase (HRP), anti-biotin antibody conjugated to HRP, LumiGLO[®] chemiluminescent reagent and peroxide.
- Prestained Protein Marker, Broad Range (Premixed Format) #7720
- Biotinylated Protein Ladder Detection Pack #7727
- Blotting Membrane:** This protocol has been optimized for nitrocellulose membranes, which CST recommends. PVDF membranes may also be used.

B Protein Blotting

A general protocol for sample preparation is described below.

- Treat cells by adding fresh media containing regulator for desired time.
- Aspirate media from cultures; wash cells with 1X PBS; aspirate.
- Lyse cells by adding 1X SDS sample buffer (100 µl per well of 6-well plate or 500 µl per plate of 10 cm diameter plate). Immediately scrape the cells off the plate and transfer the extract to a microcentrifuge tube. Keep on ice.
- Sonicate for 10–15 seconds to shear DNA and reduce sample viscosity.
- Heat a 20 µl sample to 95–100°C for 5 minutes; cool on ice.
- Microcentrifuge for 5 minutes.
- Load 20 µl onto SDS-PAGE gel (10 cm x 10 cm).

NOTE: CST recommends loading prestained molecular weight markers (#7720, 10 µl/lane) to verify electrotransfer and biotinylated protein ladder (#7727, 10 µl/lane) to determine molecular weights.

- Electrotransfer to nitrocellulose or PVDF membrane.

C Membrane Blocking and Antibody Incubations

NOTE: Volumes are for 10 cm x 10 cm (100 cm²) of membrane; for different sized membranes, adjust volumes accordingly.

- (Optional) After transfer, wash nitrocellulose membrane with 25 ml TBS for 5 minutes at room temperature.
- Incubate membrane in 25 ml of blocking buffer for 1 hour at room temperature.
- Wash three times for 5 minutes each with 15 ml of TBS/T.
- Incubate membrane and primary antibody (at the appropriate dilution) in 10 ml primary antibody dilution buffer with gentle agitation overnight at 4°C.
- Wash three times for 5 minutes each with 15 ml of TBS/T.
- Incubate membrane with HRP-conjugated secondary antibody (1:2000) and HRP-conjugated anti-biotin antibody (1:1000) to detect biotinylated protein markers in 10 ml of blocking buffer with gentle agitation for 1 hour at room temperature.
- Wash three times for 5 minutes each with 15 ml of TBS/T.

D Detection of Proteins

- Incubate membrane with 10 ml LumiGLO[®] (0.5 ml 20X LumiGLO[®], 0.5 ml 20X Peroxide and 9.0 ml Milli-Q water) with gentle agitation for 1 minute at room temperature.

NOTE: LumiGLO[®] substrate can be further diluted if signal response is too fast.

- Drain membrane of excess developing solution (do not let dry), wrap in plastic wrap and expose to x-ray film. An initial 10-second exposure should indicate the proper exposure time.

NOTE: Due to the kinetics of the detection reaction, signal is most intense immediately following LumiGLO[®] incubation and declines over the following 2 hours.