

**#9551** Store at **-20°C**

# Phospho-PTEN (Ser380) Antibody

100 µl  
 (10 Western mini-blots)



**Orders** ■ 877-616-CELL (2355)  
 orders@cellsignal.com  
**Support** ■ 877-678-TECH (8324)  
 info@cellsignal.com  
**Web** ■ www.cellsignal.com

rev. 09/28/09


This product is for *in vitro* research use only and is not intended for use in humans or animals.  
 This product is not intended for use as a therapeutic or in diagnostic procedures.

Applications	Species Cross-Reactivity*	Molecular Wt.	Source
W, IP Endogenous	H, M, R	54 kDa	Rabbit**

**Background:** PTEN (phosphatase and tensin homologue deleted on chromosome ten), also referred to as MMAC (mutated in multiple advanced cancers) phosphatase, is a tumor suppressor implicated in a wide variety of human cancers (1). PTEN encodes the 403 amino acid polypeptide originally described as a dual-specificity protein phosphatase (2). The main substrates of PTEN are inositol phospholipids generated by the activation of the phosphoinositide 3-kinase (PI3K) (3). PTEN is a major negative regulator of the PI3K/Akt signaling pathway (1,4-5). PTEN possesses a carboxy-terminal noncatalytic regulatory domain containing three phosphorylation sites (Ser380, Thr382 and Thr383), which regulates its stability and may play an important role in control of its biological activity (6,7). PTEN also regulates p53 protein levels and activity (8) and is involved in G protein coupled signaling during chemotaxis (9,10).

**Specificity/Sensitivity:** Phospho-PTEN (Ser380) Antibody detects endogenous levels of PTEN only when phosphorylated at Ser380.

**Source/Purification:** Polyclonal antibodies are produced by immunizing animals with a synthetic phospho-peptide (KLH-coupled) corresponding to residues surrounding Ser380 of human PTEN. Antibodies are purified by protein A and peptide affinity chromatography.



Western blot analysis of extracts from various cell lines, using Phospho-PTEN (Ser380) Antibody (upper) or PTEN Antibody #9552 (lower). The phospho-specificity of the antibody was confirmed by treating the membrane with calf intestinal alkaline phosphatase (CIP) after Western transfer.

**Entrez-Gene ID # 5728**  
**Swiss-Prot Acc. # P60484**

**Storage:** Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at **-20°C**.  
*Do not aliquot the antibody.*

**\*Species cross-reactivity is determined by Western blot.**

**\*\*Anti-rabbit secondary antibodies must be used to detect this antibody.**


**Recommended Antibody Dilutions:**  
 Western Blotting 1:1000  
 Immunoprecipitation 1:50

**For application specific protocols please see the web page for this product at [www.cellsignal.com](http://www.cellsignal.com).**

**Please visit [www.cellsignal.com](http://www.cellsignal.com) for a complete listing of recommended companion products.**

**Background References:**

- (1) Cantley, L.C. and Neel, B.G. (1999) *Proc. Natl. Acad. Sci. USA* 96, 4240–4245.
- (2) Myers, M.P. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94, 9052–9057.
- (3) Myers, M.P. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95, 13513–13518.
- (4) Wan, X. and Helman, L.J. (2003) *Oncogene* 22, 8205–8211.
- (5) Wu, X. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95, 15587–15591.
- (6) Vazquez, F. et al. (2000) *Mol. Cell. Biol.* 20, 5010–5018.
- (7) Torres, J. and Pulido, R. (2001) *J. Biol. Chem.* 276, 993–998.
- (8) Freeman, D.J. et al. (2003) *Cancer Cell* 3, 117–130.
- (9) Funamoto, S. et al. (2002) *Cell* 109, 611–623.
- (10) Iijima, M. and Devreotes, P. (2002) *Cell* 109, 599–610.



PTEN Signaling Pathway

**IMPORTANT: For Western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.**

**Applications Key:** W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide  
**Species Cross-Reactivity Key:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebra fish B—bovine  
**Dg—dog Pg—pig Sc—S. cerevisiae All—all species expected** Species enclosed in parentheses are predicted to react based on 100% sequence homology.