

# StemLight™ Pluripotency Antibody Kit

✓ 1 Kit  
(100 tests)




**Orders** ■ 877-616-CELL (2355)  
orders@cellsignal.com  
**Support** ■ 877-678-TECH (8324)  
info@cellsignal.com  
**Web** ■ www.cellsignal.com

rev. 01/24/12

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Products Included	Quantity	Application	Dilution	Isotype
Oct-4A (C30A3) Rabbit mAb	100 tests	IF-IC	1:200	Rabbit IgG
Sox2 (D6D9) XP <sup>®</sup> Rabbit mAb	100 tests	IF-IC	1:200	Rabbit IgG
Nanog Antibody	100 tests	IF-IC	1:200	Rabbit IgG
*SSEA4 (MC813) Mouse mAb	100 tests	IF-IC	1:200	Mouse IgG3
*TRA-1-60(S) (TRA-1-60(S)) Mouse mAb	100 tests	IF-IC	1:200	Mouse IgM
*TRA-1-81 (TRA-1-81) Mouse mAb	100 tests	IF-IC	1:200	Mouse IgM



Projected confocal z-stack of human iPS cells using TRA-1-60(S) (TRA-1-60(S)) Mouse mAb (green, lower left), TRA-1-81 (TRA-1-81) Mouse mAb (green, middle left), SSEA4 (MC813) Mouse mAb (green, upper left), Oct-4A (C30A3) Rabbit mAb (green, lower right), Sox2 (D6D9) XP<sup>®</sup> Rabbit mAb (green, middle right) and Nanog Antibody (green, upper right). Actin filaments have been labeled with DY-554 phalloidin (red). Blue pseudocolor = DRAQ5<sup>®</sup> #4084 (fluorescent DNA dye).

**Description:** The StemLight™ Pluripotency Antibody Kit contains a panel of antibodies for the detection of proteins that are specifically expressed in human pluripotent cells. The kit can be used to track the pluripotent potential of human embryonic stem (ES) or induced pluripotent (iPS) cells. The loss of these markers indicates a loss of pluripotency or differentiation of the culture. The kit components are pre-optimized for parallel use in immunofluorescent analysis. Enough reagents are provided for 50 assays based on a working volume of 100 µl.

**Specificity/Sensitivity:** Each antibody in the StemLight™ Pluripotency Antibody Kit detects endogenous levels of their respective human pluripotency marker proteins.

**Source/Purification:** Nanog Antibody was produced by immunizing animals with a synthetic peptide corresponding to amino acid sequence at the amino terminus of human nanog. Antibodies are purified by Protein A and peptide affinity chromatography. Oct-4A antibody is produced by immunizing animals with recombinant protein specific to the amino terminus of human Oct-4A. Sox2 antibody is produced by immunizing animals with a synthetic peptide corresponding to amino acid sequences at the amino terminus of human Sox2. SSEA4, TRA-1-81, and TRA-1-60(S) antibodies are produced by immunizing animals with human embryonal carcinoma 2102Ep cl.2A6 cells.

**Storage:** Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

\*Note: Due to the extracellular location of the epitope, permeabilization (detergents or alcohols) should be omitted.

For application specific protocols please see the web page for this product at [www.cellsignal.com](http://www.cellsignal.com).

Please visit [www.cellsignal.com](http://www.cellsignal.com) for a complete listing of recommended companion products.

**Background:** Pluripotency is the ability of a cell to differentiate into cell types of the three germ layers, the endoderm, ectoderm and mesoderm. It is a property shared by embryonic stem cells, embryonic carcinoma and induced pluripotent cells.

Oct-4, Sox2 and Nanog are key transcriptional regulators that are highly expressed in pluripotent cells (1). Together they form a transcriptional network that maintains cells in a pluripotent state (2,3). Over-expression of Oct-4 and Sox2 along with Klf4 and c-Myc can induce pluripotency in both mouse and human somatic cells, highlighting their roles as key regulators of the transcriptional network necessary for renewal and pluripotency (4-5). It has also been demonstrated that overexpression of Oct-4, Sox2, Nanog and Lin28 can induce pluripotency in human somatic cells (6). Upon differentiation of pluripotent cultures, expression of Oct-4, Nanog and Sox2 is downregulated.


SSEA4, TRA-1-81 and TRA-1-60 antibodies recognize antigens expressed on the cell surface of all pluripotent cells. SSEA4 recognizes a glycolipid carbohydrate epitope (7). TRA-1-60(S) and TRA-1-81 antibodies recognize different proteoglycan epitopes on variants of the same protein, podocalyxin (8). These epitopes are neurominadase sensitive and resistant, respectively. Reactivity of SSEA4, TRA-1-81 and TRA-1-60 antibodies with their respective cell surface markers are lost upon differentiation of pluripotent cells, corresponding with a loss of pluripotent potential (9).

#### Background References:


- (1) Looijenga, L.H. et al. (2003) *Cancer Res.* 63, 2244–2250.
- (2) Pesce, M. and Schöler, H.R. (2001) *Stem Cells* 19, 271–278.
- (3) Pan, G. and Thomson, J.A. (2007) *Cell. Res.* 17, 42–49.
- (4) Takahashi, K. and Yamanaka, S. (2006) *Cell* 126, 663–676.
- (5) Okita, K. et al. (2007) *Nature* 448, 313–317.
- (6) Yu, J. et al. (2007) *Science* 318, 1917–1920.
- (7) Henderson, J.K. et al. (2002) *Stem Cells* 20, 329–337.
- (8) Draper, J.S. et al. (2002) *J. Anat.* 200, 249–258.
- (9) Schopperle, W.M. and DeWolf, W.C. (2007) *Stem Cells* 25, 723–730.

Rabbit monoclonal antibody is produced under license (granting certain rights including those under U. S. Patent No. 5,675,063) from Epitomics, Inc.


DRAQ5<sup>®</sup> is a registered trademark of Biostatus Limited.




Confocal immunofluorescent analysis of NTERA-2 cells, untreated (left panel) or retinoic acid-treated (10  $\mu$ M all-trans RA for 5 days) (right panel), using TRA-1-60(S) (TRA-1-60(S)) Mouse mAb (green, upper), TRA-1-81 (TRA-1-81) Mouse mAb (green, middle) and SSEA4 (MC813) Mouse mAb (green, lower). Actin filaments have been labeled with DY-554 phalloidin (red). Blue pseudocolor = DRAQ5<sup>®</sup> #4084 (fluorescent DNA dye). Note the loss of pluripotency markers (green) as cells differentiate along the neuronal lineage with retinoic acid treatment.



Confocal immunofluorescent analysis of NTERA-2 cells, untreated (left panel) or retinoic acid-treated (10  $\mu$ M all-trans RA for 5 days) (right panel), using Neurofilament-L (C28E10) Rabbit mAb #2837 (green, upper), and a  $\beta$ -III tubulin antibody (green, lower). Actin filaments have been labeled with DY-554 phalloidin (red). Blue pseudocolor = DRAQ5<sup>®</sup> #4084 (fluorescent DNA dye). Note the appearance of neuronal markers and structures as cells differentiate along the neuronal lineage with retinoic acid treatment.



Immunofluorescent analysis of NTERA-2 cells using antibodies in the StemLight™ Pluripotency Antibody Kit.



Confocal immunofluorescent analysis of NTERA-2 cells, untreated (left panel) or retinoic acid-treated (10  $\mu$ M all-trans RA for 14 days) (right panel), using Oct-4A (C30A3) Rabbit mAb (green, upper), Sox2 (D6D9) XP™ Rabbit mAb (green, middle) and Nanog Antibody (green, lower). Actin filaments have been labeled with DY-554 phalloidin (red). Note the loss of pluripotency markers (green) as cells differentiate along the neuronal lineage with retinoic acid treatment.

# Immunofluorescence Protocol

**\*IMPORTANT:** Please refer to the **APPLICATIONS** section on the front page of the data sheet to determine **IF THIS PRODUCT** is validated and approved for the specific protocol you will be using.

## A Solutions and Reagents

**NOTE:** Prepare solutions with Milli-Q or equivalently purified water.

- 1. 10X Phosphate Buffered Saline (PBS):** To prepare 1 L add 80 g sodium chloride (NaCl), 2 g potassium chloride (KCl), 14.4 g sodium phosphate, dibasic ( $\text{Na}_2\text{HPO}_4$ ) and 2.4 g potassium phosphate, monobasic ( $\text{KH}_2\text{PO}_4$ ) to 1 L  $\text{dH}_2\text{O}$ . Adjust pH to 7.4.
- Formaldehyde, 16%, methanol free, Polysciences, Inc. (cat# 18814), use fresh, store opened vials at 4°C in dark, dilute in PBS for use.
- Xylene
- Ethanol, anhydrous denatured, histological grade, 100% and 95%
- Distilled water ( $\text{dH}_2\text{O}$ )
- Blocking Buffer:** To prepare 25 mL, add 2.5 mL 10X PBS, 1.25 mL normal serum from the same species as the secondary antibody (eg. normal goat serum, normal donkey serum) and 21.25 mL  $\text{dH}_2\text{O}$  and mix well. While stirring, add 75  $\mu\text{L}$  Triton X-100 (100%).
- Antibody Dilution Buffer:** To prepare 40 mL, add 4 mL 10X PBS to 36 mL  $\text{dH}_2\text{O}$ , mix. Add 0.4 g BSA and mix well. While stirring, add 120  $\mu\text{L}$  Triton X-100 (100%).
- 10 mM Sodium Citrate Buffer:** To prepare 1 L, add 2.94 g sodium citrate trisodium salt dihydrate ( $\text{C}_6\text{H}_5\text{Na}_3\text{O}_7 \cdot 2\text{H}_2\text{O}$ ) to 1 L  $\text{dH}_2\text{O}$ . Adjust pH to 6.0.
- 1X PBS, high salt (0.4M) (high salt PBS):** To prepare 1L, add 100 ml 10X PBS to 900 ml  $\text{dH}_2\text{O}$ . Add 23.38 g NaCl and mix.
- Fluorochrome-conjugated secondary antibody

**NOTE:** When using any primary or fluorochrome-conjugated secondary antibody for the first time, titrate the antibody to determine which dilution allows for the strongest specific signal with the least background for your sample.

- Prolong® Gold Antifade Reagent (Invitrogen, Eugene, OR, Cat# P36930)

## B Specimen Preparation

### I. Cultured Cell Lines (IF-IC)

**IMPORTANT:** Please check the **APPLICATIONS** section of the data sheet to verify that this product is validated and approved for **(IF-IC)**.

**NOTE:** Cells should be grown, treated, fixed, and stained directly in multiwell plates, chamber slides, or on coverslips.

- Rinse cells briefly in PBS.
- Aspirate PBS, cover cells to a depth of 2-3 mm with 2-4% formaldehyde in PBS.

**NOTE:** Formaldehyde is toxic, use only in fume hood.

- Allow cells to fix for 15 minutes at room temperature.
- Aspirate fixative, rinse three times in PBS for 5 minutes each.
- Methanol Permeabilization Step (if required, please refer to front page):** After formaldehyde fixation, cover cells with ice-cold 100% methanol (use enough to cover cells completely to a depth of 3-5 mm, DO NOT LET CELLS DRY), incubate cells in methanol for 10 minutes at  $-20^\circ\text{C}$ , rinse in PBS for 5 minutes.
- Proceed with Immunostaining section C.

### II. Paraffin Sections (IF-P)

**IMPORTANT:** Please check the **APPLICATIONS** section of the data sheet to verify that this product is validated and approved for **(IF-P)**.

Deparaffinization/Rehydration:

- Incubate sections in three washes of xylene for 5 minutes each.
- Incubate sections in two washes of 100% ethanol for 10 minutes each.
- Incubate sections in two washes of 95% ethanol for 10 minutes each.
- Rinse sections twice in  $\text{dH}_2\text{O}$  for 5 minutes each.

Antigen Unmasking:

- Place slides in room temperature 10 mM sodium citrate buffer pH 6.0.
- Bring slides to boiling in sodium citrate buffer using water bath or microwave, then maintain at  $95-99^\circ\text{C}$  for 10 minutes.
- Cool slides for 30 minutes on bench top.
- Rinse sections in  $\text{dH}_2\text{O}$  three times for 5 minutes each.
- Rinse sections in PBS for 5 minutes.
- Proceed with Immunostaining section C.

## III. Frozen/Cryostat Sections (IF-F)

**IMPORTANT:** Please check the **APPLICATIONS** section of the data sheet to verify that this product is validated and approved for **(IF-F)**.

**NOTE:** Fresh frozen/unfixed sections should be fixed immediately in 2-4% formaldehyde as follows to preserve signaling epitopes.

- Cover sections with 2-4% formaldehyde in PBS

**NOTE:** Formaldehyde is toxic, use only in fume hood.

- Allow sections to fix for 15 minutes at room temperature.
- Rinse slides three times in PBS for 5 minutes each.

## C Immunostaining

**NOTE:** All subsequent incubations should be carried out at room temperature unless otherwise noted in a humid light-tight box or covered dish/plate to prevent drying and fluorochrome fading.

- Block specimen in Blocking Buffer for 60 minutes.
- While blocking, prepare primary antibody by diluting as indicated on datasheet in Antibody Dilution Buffer.
- Aspirate blocking solution, apply diluted primary antibody.

**NOTE:** For double-labeling, prepare a cocktail of the primary antibodies at their appropriate dilution in Antibody Dilution Buffer.

- Incubate overnight at  $4^\circ\text{C}$ .
- Rinse three times in PBS for 5 minutes each.

**OPTION:** To decrease background stain, rinse in high salt PBS for two minutes between second and third PBS rinses. Be aware, this may reduce specific staining of some antibodies.

**NOTE:** If using primary antibodies directly conjugated with Alexa Fluor® fluorochromes, then skip to step C8.

- Incubate specimen in fluorochrome-conjugated secondary antibody diluted in Antibody Dilution Buffer for 1-2 hours at room temperature in dark.

**NOTE:** For double-labeling, prepare a cocktail of fluorochrome-conjugated secondary antibodies at their appropriate dilutions in Antibody Dilution Buffer.

- Rinse in PBS/high salt PBS as in step 5.
- Coverslip slides with Prolong® Gold Antifade Reagent or apply just enough to cover cells in multiwell plate.
- Seal slides by painting around edges of coverslips with nail polish.
- For best results examine specimens immediately using appropriate excitation wavelength. For long term storage, store slides flat at  $4^\circ\text{C}$  protected from light.